

Exercise 9.1

Q1 Write the first five terms of the sequences whose n^{th} term is $a_n = n(n + 2)$.

Answer. $a_n = n(n + 2)$

Substituting $n = 1, 2, 3, 4$, and 5 , we obtain

$$a_1 = 1(1 + 2) = 3$$

$$a_2 = 2(2 + 2) = 8$$

$$a_3 = 3(3 + 2) = 15$$

$$a_4 = 4(4 + 2) = 24$$

$$a_5 = 5(5 + 2) = 35$$

Therefore, the required terms are 3, 8, 15, 24, and 35.

Page : 180 , Block Name : Exercise 9.1

Q2 Write the first five terms of the sequences whose n^{th} term is $a_n = \frac{n}{n+1}$.

Answer. $a_n = \frac{n}{n+1}$

Substituting $n = 1, 2, 3, 4, 5$, we obtain

$$a_1 = \frac{1}{1+1} = \frac{1}{2}, a_2 = \frac{2}{2+1} = \frac{2}{3}, a_3 = \frac{3}{3+1} = \frac{3}{4}, a_4 = \frac{4}{4+1} = \frac{4}{5}, a_5 = \frac{5}{5+1} = \frac{5}{6}$$

Therefore, the required terms are $\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}$, and $\frac{5}{6}$

Page : 180 , Block Name : Exercise 9.1

Q3 Write the first five terms of the sequences whose n^{th} term is $a_n = 2^n$

Answer. $a_n = 2^n$

Substituting $n = 1, 2, 3, 4, 5$, we obtain

$$a_1 = 2^1 = 2$$

$$a_2 = 2^2 = 4$$

$$a_3 = 2^3 = 8$$

$$a_4 = 2^4 = 16$$

$$a_5 = 2^5 = 32$$

Therefore, the required terms are 2, 4, 8, 16, and 32.

Page : 180 , Block Name : Exercise 9.1

Q4 Write the first five terms of the sequences whose n^{th} term is $a_n = \frac{2n-3}{6}$

Answer. Substituting $n = 1, 2, 3, 4, 5$, we obtain

$$a_1 = \frac{2 \times 1 - 3}{6} = \frac{-1}{6}$$

$$a_2 = \frac{2 \times 2 - 3}{6} = \frac{1}{6}$$

$$a_3 = \frac{2 \times 3 - 3}{6} = \frac{3}{6} = \frac{1}{2}$$

$$a_4 = \frac{2 \times 4 - 3}{6} = \frac{5}{6}$$

$$a_5 = \frac{2 \times 5 - 3}{6} = \frac{7}{6}$$

Therefore, the required terms are $\frac{-1}{6}, \frac{1}{6}, \frac{1}{2}, \frac{5}{6}$, and $\frac{7}{6}$

Page : 180 , Block Name : Exercise 9.1

Q5 Write the first five terms of the sequences whose n^{th} term is $a_n = (-1)^{n-1}5^{n+1}$

Answer. Substituting $n = 1, 2, 3, 4, 5$, we obtain

$$a_1 = (-1)^{1-1}5^{1+1} = 5^2 = 25$$

$$a_2 = (-1)^{2-1}5^{2+1} = -5^3 = -125$$

$$a_3 = (-1)^{3-1}5^{3+1} = 5^4 = 625$$

$$a_4 = (-1)^{4-1}5^{4+1} = -5^5 = -3125$$

$$a_5 = (-1)^{5-1}5^{5+1} = 5^6 = 15625$$

Therefore, the required terms are 25, -125, 625, -3125, and 15625.

Page : 180 , Block Name : Exercise 9.1

Q6 Write the first five terms of the sequences whose n^{th} term is $a_n = n^{\frac{n^2+5}{4}}$

Answer. Substituting $n = 1, 2, 3, 4, 5$, we obtain

$$a_1 = 1 \cdot \frac{1^2 + 5}{4} = \frac{6}{4} = \frac{3}{2}$$

$$a_2 = 2 \cdot \frac{2^2 + 5}{4} = 2 \cdot \frac{9}{4} = \frac{9}{2}$$

$$a_3 = 3 \cdot \frac{3^2 + 5}{4} = 3 \cdot \frac{14}{4} = \frac{21}{2}$$

$$a_4 = 4 \cdot \frac{4^2 + 5}{4} = 21$$

$$a_5 = 5 \cdot \frac{5^2 + 5}{4} = 5 \cdot \frac{30}{4} = \frac{75}{2}$$

Therefore, the required terms are $\frac{3}{2}, \frac{9}{2}, \frac{21}{2}, 21$, and $\frac{75}{2}$.

Page : 180 , Block Name : Exercise 9.1

Q7 Find the 17th term in the following sequence whose n^{th} term is $a_n = 4n - 3$; a_{17}, a_{24}

Answer. Substituting $n = 17$, we obtain

$$a_{17} = 4(17) - 3 = 68 - 3 = 65$$

Substituting $n = 24$, we obtain

$$a_{24} = 4(24) - 3 = 96 - 3 = 93$$

Page : 180 , Block Name : Exercise 9.1

Q8 Find the 7th term in the following sequence whose n^{th} term is $a_n = \frac{n^2}{2^n}$; a_7

Answer. Substituting $n = 7$, we obtain

$$a_7 = \frac{7^2}{2 \times 7} = \frac{7}{2}$$

Here, $a_n = \frac{n^2}{2^n}$.

Substituting $n = 7$, we obtain

$$a_7 = \frac{7^2}{2^7} = \frac{49}{128}$$

Page : 180 , Block Name : Exercise 9.1

Q9 Find the 9th term in the following sequence whose n^{th} term is $a_n = (-1)^{n-1}n^3$; a_9

Answer. Substituting $n = 9$, we obtain

$$a_9 = (-1)^{9-1}(9)^3 = (9)^3 = 729$$

Page : 180 , Block Name : Exercise 9.1

Q10 $a_n = \frac{n(n-2)}{n+3}$; a_{20}

Answer. Substituting $n = 20$, we obtain

$$a_{20} = \frac{20(20-2)}{20+3} = \frac{20(18)}{23} = \frac{360}{23}$$

Q11 Write the first five terms of the following sequence and obtain the corresponding series:

$$a_1 = 3, a_n = 3a_{n-1} + 2 \text{ for all } n > 1$$

Answer.

$$a_1 = 3, a_n = 3a_{n-1} + 2 \text{ for all } n > 1$$

$$\Rightarrow a_2 = 3a_1 + 2 = 3(3) + 2 = 11$$

$$a_3 = 3a_2 + 2 = 3(11) + 2 = 35$$

$$a_4 = 3a_3 + 2 = 3(35) + 2 = 107$$

$$a_5 = 3a_4 + 2 = 3(107) + 2 = 323$$

Hence, the first five terms of the sequence are 3, 11, 35, 107, and 323.

The corresponding series is $3 + 11 + 35 + 107 + 323 + \dots$

Q12 Write the first five terms of the following sequence and obtain the corresponding series:

$$a_1 = -1, a_n = \frac{a_{n-1}}{n}, n \geq 2$$

Answer.

$$a_1 = -1, a_n = \frac{a_{n-1}}{n}, n \geq 2$$

$$\Rightarrow a_2 = \frac{a_1}{2} = \frac{-1}{2}$$

$$a_3 = \frac{a_2}{3} = \frac{-1}{6}$$

$$a_4 = \frac{a_3}{4} = \frac{-1}{24}$$

$$a_5 = \frac{a_4}{5} = \frac{-1}{120}$$

Hence, the first five terms of the sequence are

$$-1, \frac{-1}{2}, \frac{-1}{6}, \frac{-1}{24}, \text{ and } \frac{-1}{120}$$

The corresponding series is

$$(-1) + \left(\frac{-1}{2}\right) + \left(\frac{-1}{6}\right) + \left(\frac{-1}{24}\right) + \left(\frac{-1}{120}\right) + \dots$$

Q13 Write the first five terms of the following sequence and obtain the corresponding series:

$$a_1 = a_2 = 2, a_n = a_{n-1} - 1, n > 2$$

Answer.

$$a_1 = a_2 = 2, a_n = a_{n-1} - 1, n > 2$$

$$\Rightarrow a_3 = a_2 - 1 = 2 - 1 = 1$$

$$a_4 = a_3 - 1 = 1 - 1 = 0$$

$$a_5 = a_4 - 1 = 0 - 1 = -1$$

Hence, the first five terms of the sequence are 2, 2, 1, 0, and -1.

The corresponding series is $2 + 2 + 1 + 0 + (-1) + \dots$

Q14 The Fibonacci sequence is defined by

$$1 = a_1 = a_2 \text{ and } a_n = a_{n-1} + a_{n-2}, n > 2$$

Find $\frac{a_{n+1}}{a_n}$, for $n = 1, 2, 3, 4, 5$

Answer.

$$1 = a_1 = a_2$$

$$a_n = a_{n-1} + a_{n-2}, n > 2$$

$$\therefore a_3 = a_2 + a_1 = 1 + 1 = 2$$

$$a_4 = a_3 + a_2 = 2 + 1 = 3$$

$$a_5 = a_4 + a_3 = 3 + 2 = 5$$

$$a_6 = a_5 + a_4 = 5 + 3 = 8$$

$$\therefore \text{ For } n = 1, \frac{a_{n+1}}{a_n} = \frac{a_2}{a_1} = \frac{1}{1} = 1$$

$$\text{ For } n = 2, \frac{a_{n+1}}{a_n} = \frac{a_3}{a_2} = \frac{2}{1} = 2$$

$$\text{ For } n = 3, \frac{a_{n+1}}{a_n} = \frac{a_4}{a_3} = \frac{3}{2}$$

$$\text{ For } n = 4, \frac{a_{n+1}}{a_n} = \frac{a_5}{a_4} = \frac{5}{3}$$

$$\text{ For } n = 5, \frac{a_{n+1}}{a_n} = \frac{a_6}{a_5} = \frac{8}{5}$$

Exercise 9.2

Q1 Find the sum of odd integers from 1 to 2001.

Answer. The odd integers from 1 to 2001 are 1, 3, 5, ...1999, 2001.

This sequence forms an A.P.

Here, first term, $a = 1$

Common difference, $d = 2$

$$\text{ Here, } a + (n - 1)d = 2001$$

$$\Rightarrow 1 + (n - 1)(2) = 2001$$

$$\Rightarrow 2n - 2 = 2000$$

$$\Rightarrow n = 1001$$

$$S_n = \frac{n}{2}[2a + (n - 1)d]$$

$$\begin{aligned}
 \therefore S_n &= \frac{1001}{2} [2 \times 1 + (1001 - 1) \times 2] \\
 &= \frac{1001}{2} [2 + 1000 \times 2] \\
 &= \frac{1001}{2} \times 2002 \\
 &= 1001 \times 1001 \\
 &= 1002001
 \end{aligned}$$

Thus, the sum of odd numbers from 1 to 2001 is 1002001.

Page : 185 , Block Name : Exercise 9.2

Q2 Find the sum of all natural numbers lying between 100 and 1000, which are multiples of 5.

Answer. The natural numbers lying between 100 and 1000, which are multiples of 5, are 105, 110, ... 995.

Here, $a = 105$ and $d = 5$

$$a + (n - 1)d = 995$$

$$\Rightarrow 105 + (n - 1)5 = 995$$

$$\Rightarrow (n - 1)5 = 995 - 105 = 890$$

$$\Rightarrow n - 1 = 178$$

$$\Rightarrow n = 179$$

$$\begin{aligned}
 \therefore S_n &= \frac{179}{2} [2(105) + (179 - 1)(5)] \\
 &= \frac{179}{2} [2(105) + (178)(5)] \\
 &= 179[105 + (89)5] \\
 &= (179)(105 + 445) \\
 &= (179)(550) \\
 &= 98450
 \end{aligned}$$

Thus, the sum of all natural numbers lying between 100 and 1000, which are multiples of 5, is 98450.

Page : 185 , Block Name : Exercise 9.2

Q3 In an A.P., the first term is 2 and the sum of the first five terms is one-fourth of the next five terms. Show that 20th term is -112.

Answer. First term = 2

Let d be the common difference of the A.P.

Therefore, the A.P. is 2, 2 + d, 2 + 2d, 2 + 3d, ...

Sum of first five terms = 10 + 10d

Sum of next five terms = 10 + 35d

According to the given condition,

$$10 + 10d = \frac{1}{4}(10 + 35d)$$

$$\Rightarrow 40 + 40d = 10 + 35d$$

$$\Rightarrow 30 = -5d$$

$$\Rightarrow d = -6$$

$$\therefore a_{20} = a + (20 - 1)d = 2 + (19)(-6) = 2 - 114 = -112$$

Thus, the 20th term of the A.P. is -112.

Page : 185 , Block Name : Exercise 9.2

Q4 How many terms of the A.P. $-6, -\frac{11}{2}, -5, \dots$ are needed to give the sum -25?

Answer. Let the sum of n terms of the given A.P. be -25.

It is known that, $S_n = \frac{n}{2}[2a + (n - 1)d]$,

where n = number of terms, a = first term, and d = common difference Here, a = -6

$$d = -\frac{11}{2} + 6 = \frac{-11+12}{2} = \frac{1}{2}$$

Therefore, we obtain

$$-25 = \frac{n}{2} \left[2 \times (-6) + (n - 1) \left(\frac{1}{2} \right) \right]$$

$$\Rightarrow -50 = n \left[-12 + \frac{n}{2} - \frac{1}{2} \right]$$

$$\Rightarrow -50 = n \left[-\frac{25}{2} + \frac{n}{2} \right]$$

$$\Rightarrow -100 = n(-25 + n)$$

$$\Rightarrow n^2 - 55n + 100 = 0$$

$$\Rightarrow n^2 - 5n - 20(n - 5) = 0$$

$$\Rightarrow n = 20 \text{ or } 5$$

Page : 185 , Block Name : Exercise 9.2

Q5 In an A.P., if pth term is $\frac{1}{q}$ and qth term is $\frac{1}{p}$, prove that the sum of first pq terms is $\frac{1}{2}(pq + 1)$, where $p \neq q$

Answer. It is known that the general term of an A.P. is $a_n = a + (n - 1)d$

\therefore According to the given information,

$$p^{\text{th}} \text{ term} = a_p = a + (p - 1)d = \frac{1}{q} \quad \dots (1)$$

$$q^{\text{th}} \text{ term} = a_q = a + (q - 1)d = \frac{1}{p} \quad \dots (2)$$

Subtracting (2) from (1), we obtain

$$(p - 1)d - (q - 1)d = \frac{1}{q} - \frac{1}{p}$$

$$\Rightarrow (p - 1 - q + 1)d = \frac{p - q}{pq}$$

$$\Rightarrow (p - q)d = \frac{p - q}{pq}$$

$$\Rightarrow d = \frac{1}{pq}$$

Putting the value of d in (1), we obtain

$$\begin{aligned}
a + (p-1)\frac{1}{pq} &= \frac{1}{q} \\
\Rightarrow a &= \frac{1}{q} - \frac{1}{q} + \frac{1}{pq} = \frac{1}{pq} \\
\therefore S_{pq} &= \frac{pq}{2}[2a + (pq-1)d] \\
&= \frac{pq}{2}\left[\frac{2}{pq} + (pq-1)\frac{1}{pq}\right] \\
&= 1 + \frac{1}{2}(pq-1) \\
&= \frac{1}{2}pq + 1 - \frac{1}{2} = \frac{1}{2}pq + \frac{1}{2} \\
&= \frac{1}{2}(pq+1)
\end{aligned}$$

Thus, the sum of first pq terms of the A.P. is $\frac{1}{2}(pq+1)$.

Page : 185 , Block Name : Exercise 9.2

Q6 If the sum of a certain number of terms of the A.P. 25, 22, 19, ... is 116. Find the last term.

Answer. Let the sum of n terms of the given A.P. be 116.

$$S_n = \frac{n}{2}[2a + (n-1)d]$$

Here, $a = 25$ and $d = 22 - 25 = -3$

$$\therefore S_n = \frac{n}{2}[2 \times 25 + (n-1)(-3)]$$

$$\Rightarrow 116 = \frac{n}{2}[50 - 3n + 3]$$

$$\Rightarrow 232 = n(53 - 3n) = 53n - 3n^2$$

$$\Rightarrow 3n^2 - 53n + 232 = 0$$

$$\Rightarrow 3n^2 - 24n - 29n + 232 = 0$$

$$\Rightarrow 3n(n-8) - 29(n-8) = 0$$

$$\Rightarrow (n-8)(3n-29) = 0$$

$$\Rightarrow n = 8 \text{ or } n = \frac{29}{3}$$

However, n cannot be equal to $\frac{29}{3}$. Therefore, $n = 8$

$$\therefore a_8 = \text{Last term} = a + (n-1)d = 25 + (8-1)(-3)$$

$$= 25 + (7)(-3) = 25 - 21$$

$$= 4$$

Thus, the last term of the A.P. is 4.

Page : 185 , Block Name : Exercise 9.2

Q7 Find the sum to n terms of the A.P., whose k^{th} term is $5k + 1$.

Answer. It is given that the k^{th} term of the A.P. is $5k + 1$.

$$k^{th} \text{ term} = ak = a + (k-1)d$$

$$\therefore a + (k-1)d = 5k + 1$$

$$a + kd - d = 5k + 1$$

Comparing the coefficient of k, we obtain $d = 5$

$$a - d = 1$$

$$\Rightarrow a - 5 = 1$$

$$\Rightarrow a = 6$$

$$\begin{aligned} S_n &= \frac{n}{2}[2a + (n-1)d] \\ &= \frac{n}{2}[2(6) + (n-1)(5)] \\ &= \frac{n}{2}[12 + 5n - 5] \\ &= \frac{n}{2}(5n + 7) \end{aligned}$$

Page : 185 , Block Name : Exercise 9.2

Q8 If the sum of n terms of an A.P. is $(pn + qn^2)$, where p and q are constants, find the common difference.

Answer. It is known that, $S_n = \frac{n}{2}[2a + (n-1)d]$

According to the given condition,

$$\frac{n}{2}[2a + (n-1)d] = pn + qn^2$$

$$\Rightarrow \frac{n}{2}[2a + nd - d] = pn + qn^2$$

$$\Rightarrow na + n^2 \frac{d}{2} - n \cdot \frac{d}{2} = pn + qn^2$$

Comparing the coefficients of n^2 on both sides, we obtain

$$\frac{d}{2} = q$$

$$\therefore d = 2q$$

Thus, the common difference of the A.P. is $2q$.

Page : 185 , Block Name : Exercise 9.2

Q9 The sums of n terms of two arithmetic progressions are in the ratio $5n + 4 : 9n + 6$. Find the ratio of their 18^{th} terms.

Answer. Let a_1, a_2, d_1, d_2 be the first terms and the common difference of the first and second arithmetic progression respectively.

According to the given condition,

$$\frac{\text{Sum of } n \text{ terms of first A.P.}}{\text{Sum of } n \text{ terms of second A.P.}} = \frac{5n+4}{9n+6}$$

$$\Rightarrow \frac{\frac{n}{2}[2a_1 + (n-1)d_1]}{\frac{n}{2}[2a_2 + (n-1)d_2]} = \frac{5n+4}{9n+6}$$

$$\Rightarrow \frac{2a_1 + (n-1)d_1}{2a_2 + (n-1)d_2} = \frac{5n+4}{9n+6} \quad \dots (1)$$

Substituting $n = 35$ in (1), we obtain

$$\frac{2a_1 + 34d_1}{2a_2 + 34d_2} = \frac{5(35)+4}{9(35)+6}$$

$$\Rightarrow \frac{a_1 + 17d_1}{a_2 + 17d_2} = \frac{179}{321} \quad \dots (2)$$

$$\frac{18^{th} \text{ term of first A.P.}}{18^{th} \text{ term of second A.P.}} = \frac{a_1 + 17d_1}{a_2 + 17d_2} \quad \dots (3)$$

From (2) and (3), we obtain

$$\frac{18^{\text{th}} \text{ term of first A.P.}}{18^{\text{th}} \text{ term of second A.P.}} = \frac{179}{321}$$

Thus, the ratio of 18th term of both the A.P.s is 179: 321.

Page : 185 , Block Name : Exercise 9.2

Q10 If the sum of first p terms of an A.P. is equal to the sum of the first q terms, then find the sum of the first (p + q) terms.

Answer. Let a and d be the first term and the common difference of the A.P. respectively.

Here,

$$S_p = \frac{p}{2}[2a + (p - 1)d]$$

$$S_q = \frac{q}{2}[2a + (q - 1)d]$$

According to the given condition,

$$\frac{p}{2}[2a + (p - 1)d] = \frac{q}{2}[2a + (q - 1)d]$$

$$\Rightarrow p[2a + (p - 1)d] = q[2a + (q - 1)d]$$

$$\Rightarrow 2ap + pd(p - 1) = 2aq + qd(q - 1)$$

$$\Rightarrow 2a(p - q) + d[p(p - 1) - q(q - 1)] = 0$$

$$\Rightarrow 2a(p - q) + d[p^2 - p - q^2 + q] = 0$$

$$\Rightarrow 2a(p - q) + d[(p - q)(p + q) - (p - q)] = 0$$

$$\Rightarrow 2a(p - q) + d[(p - q)(p + q - 1)] = 0$$

$$\Rightarrow 2a + d(p + q - 1) = 0$$

$$\Rightarrow d = \frac{-2a}{p + q - 1} \quad \dots (1)$$

$$\therefore S_{p+q} = \frac{p+q}{2}[2a + (p + q - 1) \cdot d]$$

$$\Rightarrow S_{p+q} = \frac{p+q}{2} \left[2a + (p + q - 1) \left(\frac{-2a}{p + q - 1} \right) \right] \quad [\text{From (1)}]$$

$$= \frac{p+q}{2} [2a - 2a]$$

$$= 0$$

Thus, the sum of the first (p + q) terms of the A.P. is

Page : 185 , Block Name : Exercise 9.2

Q11 Sum of the first p, q and r terms of an A.P. are a, b and c, respectively.

Prove that $\frac{a}{p}(q - r) + \frac{b}{q}(r - p) + \frac{c}{r}(p - q) = 0$

Answer. Let a1 and d be the first term and the common difference of the A.P. respectively.

According to the given information,

$$S_p = \frac{p}{2}[2a_1 + (p-1)d] = a$$

$$\Rightarrow 2a_1 + (p-1)d = \frac{2a}{p} \quad \dots (1)$$

$$S_q = \frac{q}{2}[2a_1 + (q-1)d] = b$$

$$\Rightarrow 2a_1 + (q-1)d = \frac{2b}{q} \quad \dots (2)$$

$$S_r = \frac{r}{2}[2a_1 + (r-1)d] = c$$

$$\Rightarrow 2a_1 + (r-1)d = \frac{2c}{r} \quad \dots (3)$$

Subtracting (2) from (1), we obtain

$$p - 1d - q - 1d = 2ap - 2bq$$

$$\Rightarrow dp - 1 - q + 1 = 2aq - 2bppq$$

$$\Rightarrow dp - q = 2aq - 2bppq$$

$$\Rightarrow d = 2aq - bppqp - q \dots (4)$$

Subtracting (3) from (2), we obtain

$$(q-1)d - (r-1)d = \frac{2b}{q} - \frac{2c}{r}$$

$$\Rightarrow d(q-1-r+1) = \frac{2b}{q} - \frac{2c}{r}$$

$$\Rightarrow d(q-r) = \frac{2br-2qc}{qr}$$

$$\Rightarrow d = \frac{2(br-qc)}{qr(q-r)} \quad \dots (5)$$

Equating both the values of d obtained in (4) and (5), we obtain

$$aq - bppqp - q = br - qcqrq - r$$

$$\Rightarrow aq - bppp - q = br - qcrq - r$$

$$\Rightarrow rq - raq - bp = pp - qbr - qc$$

$$\Rightarrow raq - bpq - r = pbr - qcp - q$$

$$\Rightarrow aqr - bprq - r = bpr - cpqp - q$$

Dividing both sides by pqr, we obtain

$$\left(\frac{a}{p} - \frac{b}{q}\right)(q-r) = \left(\frac{b}{q} - \frac{c}{r}\right)(p-q)$$

$$\Rightarrow \frac{a}{p}(q-r) - \frac{b}{q}(q-r+p-q) + \frac{c}{r}(p-q) = 0$$

$$\Rightarrow \frac{a}{p}(q-r) + \frac{b}{q}(r-p) + \frac{c}{r}(p-q) = 0$$

Thus, the given result is proved.

Page : 185 , Block Name : Exercise 9.2

Q12 The ratio of the sums of m and n terms of an A.P. is $m^2 : n^2$.

Show that the ratio of m^{th} and n^{th} term is $(2m-1) : (2n-1)$.

Answer. Let a and b be the first term and the common difference of the A.P. respectively.

According to the given condition,

$$\frac{\text{Sum of } m \text{ terms}}{\text{Sum of } n \text{ terms}} = \frac{m^2}{n^2}$$

$$\Rightarrow \frac{\frac{m}{2}[2a+(m-1)d]}{\frac{n}{2}[2a+(n-1)d]} = \frac{m^2}{n^2}$$

$$\Rightarrow \frac{2a+(m-1)d}{2a+(n-1)d} = \frac{m}{n} \quad \dots (1)$$

Putting $m = 2m-1$ and $n = 2n-1$ in (1), we obtain

$$\frac{2a+(2m-2)d}{2a+(2n-2)d} = \frac{2m-1}{2n-1}$$

$$\Rightarrow \frac{a+(m-1)d}{a+(n-1)d} = \frac{2m-1}{2n-1} \quad \dots (2)$$

$$\frac{m^{\text{th}} \text{ term of A.P.}}{n^{\text{th}} \text{ term of A.P.}} = \frac{a+(m-1)d}{a+(n-1)d} \quad \dots (3)$$

From (2) and (3), we obtain

$$\frac{m^{\text{th}} \text{ term of A.P.}}{n^{\text{th}} \text{ term of A.P.}} = \frac{2m-1}{2n-1}$$

Thus, the given result is proved.

Page : 185 , Block Name : Exercise 9.2

Q13 If the sum of n terms of an A.P. is $3n^2 + 5n$ and its mth term is 164, find the value of m.

Answer. Let a and b be the first term and the common difference of the A.P. respectively.

$$a_m = a + (m - 1)d = 164 \dots (1)$$

$$\text{Sum of n terms, } S_n = \frac{n}{2} [2a + (n - 1)d]$$

Here,

$$\frac{n}{2} [2a + nd - d] = 3n^2 + 5n$$

$$\Rightarrow na + n^2 \cdot \frac{d}{2} = 3n^2 + 5n$$

Comparing the coefficient of n^2 on both sides, we obtain

$$\frac{d}{2} = 3$$

$$\Rightarrow d = 6$$

Comparing the coefficient of n on both sides, we obtain

$$a - \frac{d}{2} = 5$$

$$\Rightarrow a - 3 = 5$$

$$\Rightarrow a = 8$$

Therefore, from (1), we obtain

$$8 + (m - 1) 6 = 164$$

$$\Rightarrow (m - 1) 6 = 164 - 8 = 156$$

$$\Rightarrow m - 1 = 26$$

$$\Rightarrow m = 27$$

Thus, the value of m is 27.

Page : 185 , Block Name : Exercise 9.2

Q14 Insert five numbers between 8 and 26 such that the resulting sequence is an A.P.

Answer. Let $A_1, A_2, A_3, A_4,$ and A_5 be five numbers between 8 and 26 such that

8, $A_1, A_2, A_3, A_4, A_5, 26$ is an A.P.

Here, $a = 8, b = 26, n = 7$

Therefore, $26 = 8 + (7 - 1) d$

$$\Rightarrow 6d = 26 - 8 = 18$$

$$\Rightarrow d = 3$$

$$A_1 = a + d = 8 + 3 = 11$$

$$A_2 = a + 2d = 8 + 2 \times 3 = 8 + 6 = 14$$

$$A_3 = a + 3d = 8 + 3 \times 3 = 8 + 9 = 17$$

$$A_4 = a + 4d = 8 + 4 \times 3 = 8 + 12 = 20$$

$$A_5 = a + 5d = 8 + 5 \times 3 = 8 + 15 = 23$$

Thus, the required five numbers between 8 and 26 are 11, 14, 17, 20, and 23.

Page : 185 , Block Name : Exercise 9.2

Q15 If $\frac{a^n + b^n}{a^{n-1} + b^{n-1}}$ is the A.M. between a and b, then find the value of n.

Answer. A.M. of a and b = $\frac{a+b}{2}$

According to the given condition,

$$\frac{a+b}{2} = \frac{a^n + b^n}{a^{n-1} + b^{n-1}}$$

$$\Rightarrow (a+b)(a^{n-1} + b^{n-1}) = 2(a^n + b^n)$$

$$\Rightarrow a^n + ab^{n-1} + ba^{n-1} + b^n = 2a^n + 2b^n$$

$$\Rightarrow ab^{n-1} + a^{n-1}b = a^n + b^n$$

$$\Rightarrow ab^{n-1} - b^n = a^n - a^{n-1}b$$

$$\Rightarrow b^{n-1}(a-b) = a^{n-1}(a-b)$$

$$\Rightarrow b^{n-1} = a^{n-1}$$

$$\Rightarrow \left(\frac{a}{b}\right)^{n-1} = 1 = \left(\frac{a}{b}\right)^0$$

$$\Rightarrow n-1 = 0$$

$$\Rightarrow n = 1$$

Page : 185 , Block Name : Exercise 9.2

Q16 Between 1 and 31, m numbers have been inserted in such a way that the resulting sequence is an

A. P. and

the ratio of 7^{th} and $(m-1)^{th}$ numbers is 5 : 9. Find the value of m.

Answer. Let A_1, A_2, \dots, A_m be m numbers such that 1, $A_1, A_2, \dots, A_m, 31$ is an A.P.

Here, $a = 1, b = 31, n = m + 2$

$$\therefore 31 = 1 + (m+2-1)d$$

$$\Rightarrow 30 = (m+1)d$$

$$\Rightarrow d = \frac{30}{m+1} \quad \dots (1)$$

$$A_1 = a + d$$

$$A_2 = a + 2d$$

$$A_3 = a + 3d \dots$$

$$\therefore A_7 = a + 7d$$

$$A_{m-1} = a + (m-1)d$$

According to the given condition,

$$\frac{a+7d}{a+(m-1)d} = \frac{5}{9}$$

$$\Rightarrow \frac{1+7\left(\frac{30}{m+1}\right)}{1+(m-1)\left(\frac{30}{m+1}\right)} = \frac{5}{9} \quad [\text{From (1)}]$$

$$\Rightarrow \frac{m+1+7(30)}{m+1+30(m-1)} = \frac{5}{9}$$

$$\Rightarrow \frac{m+1+210}{m+1+30m-30} = \frac{5}{9}$$

$$\Rightarrow \frac{m+211}{31m-29} = \frac{5}{9}$$

$$\Rightarrow 9m + 1899 = 155m - 145$$

$$\Rightarrow 155m - 9m = 1899 + 145$$

$$\Rightarrow 146m = 2044$$

$$\Rightarrow m = 14$$

Thus, the value of m is 14.

Page : 185 , Block Name : Exercise 9.2

Q17 A man starts repaying a loan as first instalment of Rs. 100. If he increases the instalment by Rs 5 every month, what amount he will pay in the 30th instalment?

Answer. The first installment of the loan is Rs 100.

The second installment of the loan is Rs 105 and so on.

The amount that the man repays every month forms an A.P.

The A.P. is 100, 105, 110, ...

First term, a = 100

Common difference, d = 5

$$A_{30} = a + (30 - 1)d$$

$$= 100 + (29)(5)$$

$$= 100 + 145$$

$$= 245$$

Thus, the amount to be paid in the 30th installment is Rs 245.

Page : 186 , Block Name : Exercise 9.2

Q18 The difference between any two consecutive interior angles of a polygon is 5° .

If the smallest angle is 120° , find the number of the sides of the polygon.

Answer. The angles of the polygon will form an A.P. with common difference d as 5° and first term a as 120° .

It is known that the sum of all angles of a polygon with n sides is $180^\circ (n - 2)$.

$$\therefore S_n = 180^\circ (n - 2)$$

$$\Rightarrow \frac{n}{2} [2a + (n - 1)d] = 180^\circ (n - 2)$$

$$\Rightarrow \frac{n}{2} [240^\circ + (n - 1)5^\circ] = 180^\circ (n - 2)$$

$$\Rightarrow n[240 + (n - 1)5] = 360(n - 2)$$

$$\Rightarrow 240n + 5n^2 - 5n = 360n - 720$$

$$\begin{aligned}
&\Rightarrow 5n^2 + 235n - 360n + 720 = 0 \\
&\Rightarrow 5n^2 - 125n + 720 = 0 \\
&\Rightarrow n^2 - 25n + 144 = 0 \\
&\Rightarrow n^2 - 16n - 9n + 144 = 0 \\
&\Rightarrow n(n - 16) - 9(n - 16) = 0 \\
&\Rightarrow (n - 9)(n - 16) = 0 \\
&\Rightarrow n = 9 \text{ or } 16
\end{aligned}$$

Page : 186 , Block Name : Exercise 9.2

Exercise 9.3

Q1 Find the 20th and n^{th} terms of the G.P. $\frac{5}{2}, \frac{5}{4}, \frac{5}{8}, \dots$

Answer. The given G.P. is $\frac{5}{2}, \frac{5}{4}, \frac{5}{8}, \dots$

Here, a = First term = $\frac{5}{2}$

r = Common ratio = $\frac{\frac{5}{4}}{\frac{5}{2}} = \frac{1}{2}$

$$a_{20} = ar^{20-1} = \frac{5}{2} \left(\frac{1}{2} \right)^{19} = \frac{5}{(2)(2)^{19}} = \frac{5}{(2)^{20}}$$

$$a_n = ar^{n-1} = \frac{5}{2} \left(\frac{1}{2} \right)^{n-1} = \frac{5}{(2)(2)^{n-1}} = \frac{5}{(2)^n}$$

Page : 192 , Block Name : Exercise 9.3

Q2 Find the 12th term of a G.P. whose 8th term is 192 and the common ratio is 2.

Answer. Common ratio, r = 2

Let a be the first term of the G.P.

$$\therefore a_8 = ar^{8-1} = ar^7$$

$$\Rightarrow ar^7 = 192$$

$$a(2)^7 = 192$$

$$a(2)^7 = (2)^6 (3)$$

$$\Rightarrow a = \frac{(2)^6 \times 3}{(2)^7} = \frac{3}{2}$$

$$\therefore a_{12} = ar^{12-1} = \left(\frac{3}{2} \right) (2)^{11} = (3)(2)^{10} = 3072$$

Page : 192 , Block Name : Exercise 9.3

Q3 The 5th, 8th and 11th terms of a G.P. are p, q and s, respectively. Show that $q^2 = ps$.

Answer. Let a be the first term and r be the common ratio of the G.P.

According to the given condition,

$$a_5 = a r^{5-1} = a r^4 = p \dots (1)$$

$$a_8 = a r^{8-1} = a r^7 = q \dots (2)$$

$$a_{11} = a r^{11-1} = a r^{10} = s \dots (3)$$

Dividing equation (2) by (1), we obtain

$$\frac{ar^7}{ar^4} = \frac{q}{p}$$

$$r^3 = \frac{q}{p} \dots (4)$$

Dividing equation (3) by (2), we obtain

$$\frac{ar^{10}}{ar^7} = \frac{s}{q}$$

$$\Rightarrow r^3 = \frac{s}{q} \dots (5)$$

Equating the values of r^3 obtained in (4) and (5), we obtain

$$\frac{q}{p} = \frac{s}{q}$$

$$\Rightarrow q^2 = ps$$

Thus, the given result is proved.

Page : 192 , Block Name : Exercise 9.3

Q4 The 4^{th} term of a G.P. is square of its second term, and the first term is -3 . Determine its 7^{th} term.

Answer. Let a be the first term and r be the common ratio of the G.P.

$$\therefore a = -3$$

$$\text{It is known that, } a_n = ar^{n-1}$$

$$\therefore a_4 = ar^3 = (-3)r^3$$

$$a_2 = ar^2 = (-3)r$$

According to the given condition,

$$(-3)r^3 = [(-3)r]^2$$

$$\Rightarrow r = 9r^2$$

$$\Rightarrow r = -3$$

$$a_7 = ar^{7-1} = ar^6 = (-3)(-3)^6 = -(3)^7 = -2187$$

Thus, the seventh term of the G.P. is -2187 .

Page : 192 , Block Name : Exercise 9.3

Q5 Which term of the following sequences:

(a) $2, 2\sqrt{2}, 4, \dots$ is 128?

(b) $\sqrt{3}, 3, 3\sqrt{3}, \dots$ is 729?

(c) $\frac{1}{3}, \frac{1}{9}, \frac{1}{27}, \dots$ is $\frac{1}{19683}$?

Answer.

(a) The given sequence is $2, 2\sqrt{2}, 4, \dots$

Here, $a = 2$ and $r = \frac{2\sqrt{2}}{2} = \sqrt{2}$

Let the n th term of the given sequence be 128.

$$a_n = ar^{n-1}$$

$$\Rightarrow (2)(\sqrt{2})^{n-1} = 128$$

$$\Rightarrow (2)(2)^{\frac{n-1}{2}} = (2)^7$$

$$\Rightarrow (2)^{\frac{n-1}{2}+1} = (2)^7$$

$$\therefore \frac{n-1}{2} = 6$$

$$\Rightarrow n - 1 = 12$$

$$\Rightarrow n = 13$$

Thus, the 13th term of the given sequence is 128.

(b) The given sequence is $\sqrt{3}, 3, 3\sqrt{3}$

Here, $a = \sqrt{3}$ and $r = \frac{3}{\sqrt{3}} = \sqrt{3}$

Let the n th term of the given sequence be 729.

$$a_n = ar^{n-1}$$

$$\therefore ar^{n-1} = 729$$

$$\Rightarrow (\sqrt{3})(\sqrt{3})^{n-1} = 729$$

$$\Rightarrow (3)^{\frac{1}{2}}(3)^{\frac{n-1}{2}} = (3)^6$$

$$\Rightarrow (3)^{\frac{1}{2} + \frac{n-1}{2}} = (3)^6$$

$$\therefore \frac{1}{2} + \frac{n-1}{2} = 6$$

$$\Rightarrow \frac{1+n-1}{2} = 6$$

$$\Rightarrow n = 12$$

Thus, the 12th term of the given sequence is 729.

(c) The given sequence is $\frac{1}{3}, \frac{1}{9}, \frac{1}{27}, \dots$

Here, $a = \frac{1}{3}$ and $r = \frac{1}{9} \div \frac{1}{3} = \frac{1}{3}$

Let the n th term of the given sequence be $\frac{1}{19683}$.

$$a_n = ar^{n-1}$$

$$\therefore ar^{n-1} = \frac{1}{19683}$$

$$\Rightarrow \left(\frac{1}{3}\right)\left(\frac{1}{3}\right)^{n-1} = \frac{1}{19683}$$

$$\Rightarrow \left(\frac{1}{3}\right)^n = \left(\frac{1}{3}\right)^9$$

$$\Rightarrow n = 9$$

Thus, the 9th term of the given sequence is $\frac{1}{19683}$.

Page : 192 , Block Name : Exercise 9.3

Q6 For what values of x , the numbers $-\frac{2}{7}, x, -\frac{7}{2}a$ are in G.P.?

Answer. The given numbers are $-\frac{2}{7}, x, -\frac{7}{2}a$.

$$\text{Common ratio} = \frac{\frac{x}{-\frac{2}{7}}}{-\frac{2}{7}} = \frac{-7x}{2}$$

Also, common ratio = $\frac{-7}{x} = \frac{-7}{2x}$

$$\therefore \frac{-7x}{2} = \frac{-7}{2x}$$

$$\Rightarrow x^2 = \frac{-2 \times 7}{-2 \times 7} = 1$$

$$\Rightarrow x = \sqrt{1}$$

$$\Rightarrow x = \pm 1$$

Thus, for $x = \pm 1$, the given numbers will be in G.P.

Page : 192 , Block Name : Exercise 9.3

Q7 Find the sum to n terms in the geometric progression 0.15, 0.015, 0.0015, ... 20 terms.

Answer. The given G.P. is 0.15, 0.015, 0.0015, ...

Here, $a = 0.15$ and $r = \frac{0.015}{0.15} = 0.1$

$$\begin{aligned} S_n &= \frac{a(1-r^n)}{1-r} \\ \therefore S_{20} &= \frac{0.15[1-(0.1)^{20}]}{1-0.1} \\ &= \frac{0.15}{0.9}[1-(0.1)^{20}] \\ &= \frac{15}{90}[1-(0.1)^{20}] \\ &= \frac{1}{6}[1-(0.1)^{20}] \end{aligned}$$

Page : 192 , Block Name : Exercise 9.3

Q8 Find the sum to n terms in the geometric progression $\sqrt{7}, \sqrt{21}, 3\sqrt{7}, \dots, n$

Answer.

The given G.P. is $\sqrt{7}, \sqrt{21}, 3\sqrt{7}, \dots$

Here, $a = \sqrt{7}$

$$r = \frac{\sqrt{21}}{\sqrt{7}} = \sqrt{3}$$

$$S_n = \frac{a(1-r^n)}{1-r}$$

$$\begin{aligned} \therefore S_n &= \frac{\sqrt{7}[1-(\sqrt{3})^n]}{1-\sqrt{3}} \\ &= \frac{\sqrt{7}[1-(\sqrt{3})^n]}{1-\sqrt{3}} \times \frac{1+\sqrt{3}}{1+\sqrt{3}} \quad [\text{by rationalizing}] \end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{7}(1+\sqrt{3})[1-(\sqrt{3})^n]}{1-3} \\
&= \frac{-\sqrt{7}(1+\sqrt{3})}{2} \left[1-(3)^{\frac{n}{2}}\right] \\
&= \frac{\sqrt{7}(1+\sqrt{3})}{2} \left[(3)^{\frac{n}{2}}-1\right]
\end{aligned}$$

Page : 192 , Block Name : Exercise 9.3

Q9

Find the sum to n terms in the geometric progression

$1, -a, a^2, -a^3 \dots$ (if $a \neq -1$)

Answer.

The given G.P. is $1, -a, a^2, -a^3, \dots$

Here, first term $= a_1 = 1$

Common ratio $= r = -a$

$$S_n = \frac{a_1(1-r^n)}{1-r}$$

$$\therefore S_n = \frac{1[1-(-a)^n]}{1-(-a)} = \frac{[1-(-a)^n]}{1+a}$$

Page : 192 , Block Name : Exercise 9.3

Q10 Find the sum to n terms in the geometric progression $x^3, x^5, x^7, \dots n$ terms (if $x \neq \pm 1$)

Answer.

The given G.P. is x^3, x^5, x^7, \dots

Here, $a = x^3$ and $r = x^2$

$$S_n = \frac{a(1-r^n)}{1-r} = \frac{x^3[1-(x^2)^n]}{1-x^2} = \frac{x^3(1-x^{2n})}{1-x^2}$$

Page : 192 , Block Name : Exercise 9.3

Q11 Evaluate $\sum_{k=1}^{11} (2 + 3^k)$

Answer.

$$\sum_{k=1}^{11} (2 + 3^k) = \sum_{k=1}^{11} (2) + \sum_{k=1}^{11} 3^k = 2(11) + \sum_{k=1}^{11} 3^k = 22 + \sum_{k=1}^{11} 3^k \quad \dots (1)$$

$$\sum_{k=1}^{11} 3^k = 3^1 + 3^2 + 3^3 + \dots + 3^{11}$$

The terms of this sequence $3, 3^2, 3^3, \dots$ forms a G.P.

$$S_n = \frac{a(r^n-1)}{r-1}$$

$$\Rightarrow S_n = \frac{3[(3)^{11}-1]}{3-1}$$

$$\Rightarrow S_n = \frac{3}{2} (3^{11} - 1)$$

$$\therefore \sum_{k=1}^{11} 3^k = \frac{3}{2} (3^{11} - 1)$$

Substituting this value in equation (1), we obtain

$$\sum_{k=1}^{11} (2 + 3^k) = 22 + \frac{3}{2} (3^{11} - 1)$$

Page : 192 , Block Name : Exercise 9.3

Q12 The sum of first three terms of a G.P. is $\frac{39}{10}$ and their product is 1.
Find the common ratio and the terms.

Answer.

Let r, a, ar be the first three terms of the G.P.

$$\frac{a}{r} + a + ar = \frac{39}{10} \quad \dots (1)$$

$$\left(\frac{a}{r}\right)(a)(ar) = 1 \quad \dots (2)$$

From (2), we obtain

$$a^3 = 1$$

$$\Rightarrow a = 1 \text{ (Considering real roots only)}$$

Substituting $a = 1$ in equation (1), we obtain

$$\frac{1}{r} + 1 + r = \frac{39}{10}$$

$$\Rightarrow 1 + r + r^2 = \frac{39}{10}r$$

$$\Rightarrow 10 + 10r + 10r^2 - 39r = 0$$

$$\Rightarrow 10r^2 - 29r + 10 = 0$$

$$\Rightarrow 10r^2 - 25r - 4r + 10 = 0$$

$$\Rightarrow 5r(2r - 5) - 2(2r - 5) = 0$$

$$\Rightarrow (5r - 2)(2r - 5) = 0$$

$$\Rightarrow r = \frac{2}{5} \text{ or } \frac{5}{2}$$

Thus, the three terms of G.P. are $\frac{5}{2}, 1$, and $\frac{2}{5}$

Page : 192 , Block Name : Exercise 9.3

Q13 How many terms of G.P. $3, 3^2, 3^3, \dots$ are needed to give the sum 120?

Answer. The given G.P. is $3, 3^2, 3^3, \dots$

Let n terms of this G.P. be required to obtain the sum as 120.

$$S_n = \frac{a(r^n - 1)}{r - 1}$$

Here, $a = 3$ and $r = 3$

$$\therefore S_n = 120 = \frac{3(3^n - 1)}{3 - 1}$$

$$\Rightarrow 120 = \frac{3(3^n - 1)}{2}$$

$$\Rightarrow \frac{120 \times 2}{3} = 3^n - 1$$

$$\Rightarrow 3^n - 1 = 80$$

$$\Rightarrow 3^n = 81$$

$$\therefore n = 4$$

Thus, four terms of the given G.P. are required to obtain the sum as 120.

Page : 192 , Block Name : Exercise 9.3

Q14 The sum of first three terms of a G.P. is 16 and the sum of the next three terms is 128.
Determine the first term, the common ratio and the sum to n terms of the G.P.

Answer.

Let the G.P. be a, ar, ar^2, ar^3, \dots

According to the given condition,

$$a + ar + ar^2 = 16 \text{ and } ar^3 + ar^4 + ar^5 = 128$$

$$\Rightarrow a(1 + r + r^2) = 16 \dots (1)$$

$$ar^3(1 + r + r^2) = 128 \dots (2)$$

Dividing equation (2) by (1), we obtain

$$\frac{ar^3(1+r+r^2)}{a(1+r+r^2)} = \frac{128}{16}$$

$$\Rightarrow r^3 = 8$$

$$\therefore r = 2$$

Substituting $r = 2$ in (1), we obtain

$$a(1 + 2 + 4) = 16$$

$$\Rightarrow a(7) = 16$$

$$\Rightarrow a = \frac{16}{7}$$

$$S_n = \frac{a(r^n - 1)}{r - 1}$$

$$\Rightarrow S_n = \frac{16}{7} \frac{(2^n - 1)}{2 - 1} = \frac{16}{7}(2^n - 1)$$

Page : 192 , Block Name : Exercise 9.3

Q15 Given a G.P. with $a = 729$ and 7th term 64, determine S_7 .

Answer. $a = 729$

$$a_7 = 64$$

Let r be the common ratio of the G.P.

It is known that, $a_n = ar^{n-1}$

$$a_7 = ar^{7-1} = (729)r^6$$

$$\Rightarrow 64 = 729r^6$$

$$\Rightarrow r^6 = \frac{64}{729}$$

$$\Rightarrow r^6 = \left(\frac{2}{3}\right)^6$$

$$\Rightarrow r = \frac{2}{3}$$

Also, it is known that, $S_n = \frac{a(1-r^n)}{1-r}$

$$\begin{aligned}\therefore S_7 &= \frac{729 \left[1 - \left(\frac{2}{3}\right)^7 \right]}{1 - \frac{2}{3}} \\ &= 3 \times 729 \left[1 - \left(\frac{2}{3}\right)^7 \right] \\ &= (3)^7 - \left(\frac{(3)^7 - (2)^7}{(3)^7} \right) \\ &= 2187 - 128 \\ &= 2059\end{aligned}$$

Page : 192 , Block Name : Exercise 9.3

Q16 Find a G.P. for which sum of the first two terms is -4 and the fifth term is 4 times the third term.

Answer. Let a be the first term and r be the common ratio of the G.P.

According to the given conditions,

$$S_2 = -4 = \frac{a(1-r^2)}{1-r} \quad \dots (1)$$

$$a_5 = 4 \times a_3$$

$$a_5^4 = 4ar^2$$

$$\Rightarrow r^2 = 4$$

$$\therefore r = \pm 2$$

From (1), we obtain

$$-4 = \frac{a[1-(2)^2]}{1-2} \text{ for } r = 2$$

$$\Rightarrow -4 = \frac{a(1-4)}{-1}$$

$$\Rightarrow -4 = a(3)$$

$$\Rightarrow a = \frac{-4}{3}$$

$$\text{Also, } -4 = \frac{a[1-(-2)^2]}{1-(-2)} \text{ for } r = -2$$

$$\Rightarrow -4 = \frac{a(1-4)}{1+2}$$

$$\Rightarrow -4 = \frac{a(-3)}{3}$$

$$\Rightarrow a = 4$$

Thus, the required G.P. is

$$\frac{-4}{3}, \frac{-8}{3}, \frac{-16}{3}, \dots, 4, -8, 16, -32, \dots$$

Page : 192 , Block Name : Exercise 9.3

Q17 If the 4^{th} , 10^{th} and 16^{th} terms of a G.P. are x , y and z , respectively. Prove that x , y , z are in G.P.

Answer. Let a be the first term and r be the common ratio of the G.P.

According to the given condition,

$$a_4 = ar^3 = x \dots (1)$$

$$a_{10} = ar^9 = y \dots (2)$$

$$a_{16} = ar^{15} = z \dots (3)$$

Dividing (2) by (1), we obtain

$$\frac{y}{x} = \frac{ar^9}{ar^3} \Rightarrow \frac{y}{x} = r^6$$

Dividing (3) by (2), we obtain

$$\frac{z}{y} = \frac{ar^{15}}{ar^9} \Rightarrow \frac{z}{y} = r^6$$

$$\therefore \frac{y}{x} = \frac{z}{y}$$

Thus, x , y , z are in G. P.

Page : 192 , Block Name : Exercise 9.3

Q18 Find the sum to n terms of the sequence, 8, 88, 888, 8888... .

Answer. The given sequence is 8, 88, 888, 8888...

This sequence is not a G.P. However, it can be changed to G.P. by writing the terms as

$S_n = 8 + 88 + 888 + 8888 + \dots$ to n terms

$$= \frac{8}{9} [9 + 99 + 999 + 9999 + \dots \text{ to } n \text{ terms}]$$

$$= \frac{8}{9} [(10 - 1) + (10^2 - 1) + (10^3 - 1) + (10^4 - 1) + \dots \text{ to } n \text{ terms}]$$

$$= \frac{8}{9} [(10 + 10^2 + \dots \text{ } n \text{ terms}) - (1 + 1 + 1 + \dots \text{ } n \text{ terms})]$$

$$= \frac{8}{9} \left[\frac{10(10^n - 1)}{10 - 1} - n \right]$$

$$= \frac{8}{9} \left[\frac{10(10^n - 1)}{9} - n \right]$$

$$= \frac{80}{81} (10^n - 1) - \frac{8}{9} n$$

Page : 193 , Block Name : Exercise 9.3

Q19 Find the sum of the products of the corresponding terms of the sequences 2, 4, 8, 16, 32 and 128, 32, 8, 2, $\frac{1}{2}$.

Answer.

$$\begin{aligned}\text{Required sum} &= 2 \times 128 + 4 \times 32 + 8 \times 8 + 16 \times 2 + 32 \times \frac{1}{2} \\ &= 64 \left[4 + 2 + 1 + \frac{1}{2} + \frac{1}{2^2} \right]\end{aligned}$$

Here, $4, 2, 1, \frac{1}{2}, \frac{1}{2^2}$ is a G.P.

First term, $a = 4$

Common ratio, $r = \frac{1}{2}$

$$S_n = \frac{a(1-r^n)}{1-r}$$

$$\therefore S_5 = \frac{4 \left[1 - \left(\frac{1}{2} \right)^5 \right]}{1 - \frac{1}{2}} = \frac{4 \left[1 - \frac{1}{32} \right]}{\frac{1}{2}} = 8 \left(\frac{32-1}{32} \right) = \frac{31}{4}$$

$$\therefore \text{Required sum} = 64 \left(\frac{31}{4} \right) = (16)(31) = 496$$

Page : 193 , Block Name : Exercise 9.3

Q20 Show that the products of the corresponding terms of the sequences $a, ar, ar^2 \dots ar^{n-1}$ and $A, AR, AR^2, \dots AR^{n-1}$ form a G.P. and find the common ratio.

Answer. It has to be proved that the sequence, $a, ar, ar^2 \dots ar^{n-1}$ and $A, AR, AR^2, \dots AR^{n-1}$ form a G.P.

$$\frac{\text{Second term}}{\text{First term}} = \frac{arAR}{aA} = rR$$

$$\frac{\text{Third term}}{\text{Second term}} = \frac{ar^2AR^2}{arAR} = rR$$

Thus, the above sequence forms a G.P. and the common ratio is rR .

Page : 193 , Block Name : Exercise 9.3

Q21 Find four numbers forming a geometric progression in which the third term is greater than the first term by 9,
and the second term is greater than the 4th by 18.

Answer. Let a be the first term and r be the common ratio of the G.P.

$$a_1 = a, a_2 = ar, a_3 = ar^2, a_4 = ar^3$$

By the given condition,

$$a_3 = a_1 + 9$$

$$\Rightarrow ar^2 = a + 9 \dots (1)$$

$$a_2 = a_4 + 18$$

$$\Rightarrow ar = ar^3 + 18 \dots (2)$$

From (1) and (2), we obtain

$$a(r^2 - 1) = 9 \dots (3)$$

$$ar(1 - r^2) = 18 \dots (4)$$

Dividing (4) by (3), we obtain

$$\frac{ar(1-r^2)}{a(r^2-1)} = \frac{18}{9}$$

$$\Rightarrow -r = 2$$

$$\Rightarrow r = -2$$

Substituting the value of r in (1), we obtain

$$4a = a + 9$$

$$\Rightarrow 3a = 9$$

$$\therefore a = 3$$

Thus, the first four numbers of the G.P. are 3, 3(-2), 3(-2)2, and 3(-2)3 i.e., 3, -6, 12, and -24.

Page : 193 , Block Name : Exercise 9.3

Q22 If the p^{th} , q^{th} and r^{th} terms of a G.P. are a, b and c, respectively.

Prove that $a^{q-r}b^{r-p}c^{p-q} = 1$

Answer. Let A be the first term and R be the common ratio of the G.P.

According to the given information,

$$AR^{p-1} = a$$

$$AR^{q-1} = b$$

$$AR^{r-1} = c$$

$$a^{q-r}b^{r-p}c^{p-q}$$

$$= A^{q-r} \times R^{(p-1)(q-r)} \times A^{r-p} \times R^{(q-1)(r-p)} \times A^{p-q} \times R^{(r-1)(p-q)}$$

$$= Aq^{-r+r-p+p-q} \times R(pr - pr - q + r) + (rq - r + p - pq) + (pr - p - qr + q)$$

$$= A^0 \times R^0$$

$$= 1$$

Thus, the given result is proved.

Page : 193 , Block Name : Exercise 9.3

Q23 If the first and the n^{th} term of a G.P. are a and b, respectively, and if P is the product of n terms, prove that $P^2 = (ab)^n$.

Answer. The first term of the G.P is a and the last term is b.

Therefore, the G.P. is $a, ar^2, ar^3, \dots, ar^{n-1}$ where r is the common ratio.

$$b = ar^{n-1} \dots (1)$$

P = Product of n terms

$$= (a)(ar) (ar^2) \dots (ar^{n-1})$$

$$= (a \times a \times \dots a) (r \times r^2 \times \dots r^{n-1})$$

$$= a^n r^{1+2+\dots+(n-1)} \dots (2)$$

Here, 1, 2, ..., (n - 1) is an A.P.

$$\therefore 1 + 2 + \dots + (n - 1)$$

$$= \frac{n-1}{2} [2 + (n - 1 - 1) \times 1] = \frac{n-1}{2} [2 + n - 2] = \frac{n(n-1)}{2}$$

$$P = a^n r^{\frac{n(n-1)}{2}}$$

$$\therefore P^2 = a^{2n} r^{n(n-1)}$$

$$\begin{aligned}
&= \left[a^2 r^{(n-1)} \right]^n \\
&= \left[a \times ar^{n-1} \right]^n \\
&= (ab)^n \quad [\text{using (1)}]
\end{aligned}$$

Thus, the given result is proved.

Page : 193 , Block Name : Exercise 9.3

Q24 Show that the ratio of the sum of first n terms of a G.P. to the sum of terms from $(n+1)^{\text{th}}$ to $(2n)^{\text{th}}$ term is $\frac{1}{r^n}$

Answer. Let a be the first term and r be the common ratio of the G.P.

$$\text{Sum of 1st } n \text{ terms} = \frac{a(1-r^n)}{(1-r)}$$

$$\text{Since there are } n \text{ terms from } (n+1)^{\text{th}} \text{ to } (2n)^{\text{th}} \text{ term,} = \frac{a_{n+1}(1-r^n)}{(1-r)}$$

$$\text{Thus, required ratio} = \frac{a(1-r^n)}{(1-r)} \times \frac{(1-r)}{ar^n(1-r^n)} = \frac{1}{r^n}$$

Thus, the ratio of the sum of first n terms of a G.P. to the sum of terms from $(n+1)^{\text{th}}$ to $(2n)^{\text{th}}$ term is $\frac{1}{r^n}$.

Page : 193 , Block Name : Exercise 9.3

Q25 If a, b, c and d are in G.P. show that

$$(a^2 + b^2 + c^2)(b^2 + c^2 + d^2) = (ab + bc + cd)^2$$

Answer. a, b, c, d are in G.P. Therefore,

$$bc = ad \dots (1)$$

$$b^2 = ac \dots (2)$$

$$c^2 = bd \dots (3)$$

It has to be proved that,

$$(a^2 + b^2 + c^2)(b^2 + c^2 + d^2) = (ab + bc + cd)^2$$

R.H.S.

$$= (ab + bc + cd)^2$$

$$= (ab + ad + cd)^2 [U \text{ sing (1)}]$$

$$= [ab + d(a + c)]^2$$

$$= a^2b^2 + 2abd(a + c) + d^2(a + c)^2$$

$$= a^2b^2 + 2a^2c^2 + 2acbd + d^2(a^2 + 2ac + c^2)$$

$$= a^2b^2 + 2a^2c^2 + 2b^2c^2 + d^2a^2 + 2a^2b^2 + d^2c^2 \quad [U \text{ sing (1) and (2)}]$$

$$= a^2b^2 + a^2c^2 + a^2c^2 + b^2c^2 + b^2c^2 + d^2a^2 + d^2b^2 + d^2b^2$$

$$+ d^2c^2$$

$$= a^2b^2 + a^2c^2 + a^2d^2 + b^2 \times b^2 + b^2c^2 + b^2d^2 + c^2b^2 + c^2 \times c^2 + c^2d^2$$

[Using (2) and (3) and rearranging terms]

$$\begin{aligned}
&= a^2 (b^2 + c^2 + d^2) + b^2 (b^2 + c^2 + d^2) + c^2 (b^2 + c^2 + d^2) \\
&= (a^2 + b^2 + c^2) (b^2 + c^2 + d^2) \\
&= \text{L.H.S.} \\
&\therefore \text{L.H.S.} = \text{R.H.S.} \\
&\therefore (a^2 + b^2 + c^2) (b^2 + c^2 + d^2) = (ab + bc + cd)^2
\end{aligned}$$

Page : 193 , Block Name : Exercise 9.3

Q26 Insert two numbers between 3 and 81 so that the resulting sequence is G.P.

Answer.

Let G_1 and G_2 be two numbers between 3 and 81 such that the series, 3, G_1 , G_2 , 81, forms a G.P.

Let a be the first term and r be the common ratio of the G.P.

$$\therefore 81 = (3)(r)^3$$

$$\Rightarrow r^3 = 27$$

$$\therefore r = 3 \text{ (Taking real roots only)}$$

For $r = 3$

$$G_1 = ar = (3)(3) = 9$$

$$G_2 = ar^2 = (3)(3)^2 = 27$$

Thus, the required two numbers are 9 and 27.

Page : 193 , Block Name : Exercise 9.3

Q27 Find the value of n so that $\frac{a^{n+1}+b^{n+1}}{a^n+b^n}$ may be the geometric mean between a and b .

Answer.

G. M. of a and b is \sqrt{ab}

By the given condition, $\frac{a^{n+1}+b^{n+1}}{a^n+b^n} = \sqrt{ab}$

Squaring both sides, we obtain

$$\begin{aligned}
\frac{(a^{n+1}+b^{n+1})^2}{(a^n+b^n)^2} &= ab \\
\Rightarrow a^{2n+2} + 2a^{n+1}b^{n+1} + b^{2n+2} &= (ab)(a^{2n} + 2a^n b^n + b^{2n}) \\
\Rightarrow a^{2n+2} + 2a^{n+1}b^{n+1} + b^{2n+2} &= ab^{2n+1} + 2a^{2n+1}b + ab^{2n+1} \\
\Rightarrow a^{2n+2} + b^{2n+2} + 2a^{n+1}b^{n+1} &= ab^{2n+1} + 2a^{2n+1}b + ab^{2n+1} \\
\Rightarrow a^{2n+2} - a^{2n+1}b &= ab^{2n+1} - b^{2n+2} \\
\Rightarrow a^{2n+1}(a - b) &= b^{2n+1}(a - b) \\
\Rightarrow \left(\frac{a}{b}\right)^{2n+1} &= 1 = \left(\frac{a}{b}\right)^0 \\
\Rightarrow 2n + 1 &= 0 \\
\Rightarrow n &= \frac{-1}{2}
\end{aligned}$$

Page : 193 , Block Name : Exercise 9.3

Q28 The sum of two numbers is 6 times their geometric mean, show that numbers are in the ratio $(3 + 2\sqrt{2}) : (3 - 2\sqrt{2})$

Answer. Let the two numbers be a and b.

$$G.M. = \sqrt{ab}$$

According to the given condition,

$$a + b = 6\sqrt{ab}$$

$$\Rightarrow (a + b)^2 = 36(ab) \quad \dots (1)$$

Also,

$$(a - b)^2 = (a + b)^2 - 4ab = 36ab - 4ab = 32ab$$

$$\Rightarrow a - b = \sqrt{32}\sqrt{ab}$$

$$= 4\sqrt{2}\sqrt{ab} \quad \dots (2)$$

Adding (1) and (2), we obtain

$$2a = (6 + 4\sqrt{2})\sqrt{ab}$$

$$\Rightarrow a = (3 + 2\sqrt{2})\sqrt{ab}$$

Substituting the value of a in (1), we obtain

$$b = 6\sqrt{ab} - (3 + 2\sqrt{2})\sqrt{ab}$$

$$\Rightarrow b = (3 - 2\sqrt{2})\sqrt{ab}$$

$$\frac{a}{b} = \frac{(3+2\sqrt{2})\sqrt{ab}}{(3-2\sqrt{2})\sqrt{ab}} = \frac{3+2\sqrt{2}}{3-2\sqrt{2}}$$

Thus, the required ratio is $(3 + 2\sqrt{2}) : (3 - 2\sqrt{2})$

Page : 193 , Block Name : Exercise 9.3

Q29 If A and G be A.M. and G.M., respectively between two positive numbers, prove that the numbers are $A \pm \sqrt{(A + G)(A - G)}$.

Answer. It is given that A and G are A.M. and G.M. between two positive numbers.

Let these two positive numbers be a and b.

$$\therefore AM = A = \frac{a+b}{2} \quad \dots (1)$$

$$GM = G = \sqrt{ab} \quad \dots (2)$$

From (1) and (2), we obtain

$$a + b = 2A \quad \dots (3)$$

$$ab = G^2 \quad \dots (4)$$

Substituting the value of a and b from (3) and (4) in the

identity $(a - b)^2 = (a + b)^2 - 4ab$, we obtain

$$(a - b)^2 = 4A^2 - 4G^2 = 4(A^2 - G^2)$$

$$(a - b)^2 = 4(A + G)(A - G)$$

$$(a - b) = 2\sqrt{(A + G)(A - G)} \quad \dots (5)$$

From (3) and (5), we obtain

$$2a = 2A + 2\sqrt{(A+G)(A-G)}$$

$$\Rightarrow a = A + \sqrt{(A+G)(A-G)}$$

Substituting the value of a in (3), we obtain

$$b = 2A - A - \sqrt{(A+G)(A-G)} = A - \sqrt{(A+G)(A-G)}$$

Thus, the two numbers are $A \pm \sqrt{(A+G)(A-G)}$

Page : 193 , Block Name : Exercise 9.3

Q30 The number of bacteria in a certain culture doubles every hour. If there were 30 bacteria present in the culture originally,
how many bacteria will be present at the end of 2nd hour, 4th hour and nth hour ?

Answer. It is given that the number of bacteria doubles every hour. Therefore, the number of bacteria after every hour will form a G.P.

Here, $a = 30$ and $r = 2$

Here, $a = 30$ and $r = 2$

$$\therefore a_3 = ar^2 = (30)(2)^2 = 120$$

Therefore, the number of bacteria at the end of 2nd hour will be 120.

The number of bacteria at the end of 4th hour will be 480.

$$a_{n+1} = ar^n = (30)2^n$$

Thus, number of bacteria at the end of nth hour will be

$$30(2)^n.$$

Page : 193 , Block Name : Exercise 9.3

Q31 What will Rs 500 amounts to in 10 years after its deposit in a bank which pays annual interest rate of 10% compounded annually?

Answer. The amount deposited in the bank is Rs 500.

At the end of first year, amount = Rs 500 $\left(1 + \frac{1}{10}\right)$ = Rs 500 (1.1)

At the end of 2nd year, amount = Rs 500 (1.1) (1.1)

At the end of 3rd year, amount = Rs 500 (1.1) (1.1) (1.1) and so on

\therefore Amount at the end of 10 years = Rs 500 (1.1) (1.1) ... (10 times)

$$= \text{Rs } 500(1.1)^{10}$$

Page : 193 , Block Name : Exercise 9.3

Q32 If A.M. and G.M. of roots of a quadratic equation are 8 and 5, respectively, then obtain the quadratic equation.

Answer. Let the root of the quadratic equation be a and b.

According to the given condition,

$$A \cdot M. = \frac{a+b}{2} = 8 \Rightarrow a+b = 16 \quad \dots (1)$$

$$G.M = \sqrt{ab} = 5 \Rightarrow ab = 25 \quad \dots (2)$$

The quadratic equation is given by,

$$x^2 - x(\text{sum of roots}) + (\text{Product of roots}) = 0$$

$$x^2 - x(a+b) + (\text{Product of roots}) = 0$$

$$x^2 - 16x + 25 = 0 \text{ [using (1) and (2)]}$$

Thus, the required quadratic equation is $x^2 - 16x + 25 = 0$.

Page : 193 , Block Name : Exercise 9.3

Exercise 9.4

Q1 Find the sum to n terms of the series $1 \times 2 + 2 \times 3 + 3 \times 4 + 4 \times 5 + \dots$

Answer. The given series is $1 \times 2 + 2 \times 3 + 3 \times 4 + 4 \times 5 + \dots$

nth term, $a_n = n(n+1)$

$$\begin{aligned} \therefore S_n &= \sum_{k=1}^n a_k = \sum_{k=1}^n k(k+1) \\ &= \sum_{k=1}^n k^2 + \sum_{k=1}^n k \\ &= \frac{n(n+1)(2n+1)}{6} + \frac{n(n+1)}{2} \\ &= \frac{n(n+1)}{2} \left(\frac{2n+1}{3} + 1 \right) \\ &= \frac{n(n+1)}{2} \left(\frac{2n+1+3}{3} \right) \\ &= \frac{n(n+1)(n+2)}{3} \end{aligned}$$

Page : 196 , Block Name : Exercise 9.4

Q2 Find the sum to n terms of the series $1 \times 2 \times 3 + 2 \times 3 \times 4 + 3 \times 4 \times 5 + \dots$

Answer. The given series is $1 \times 2 \times 3 + 2 \times 3 \times 4 + 3 \times 4 \times 5 + \dots$

nth term, $a_n = n(n+1)(n+2)$

$$\begin{aligned}
&= (n^2 + n)(n + 2) \\
&= n^3 + 3n^2 + 2n \\
\therefore S_n &= \sum_{k=1}^n a_k \\
&= \sum_{k=1}^n k^3 + 3 \sum_{k=1}^n k^2 + 2 \sum_{k=1}^n k \\
&= \left[\frac{n(n+1)}{2} \right]^2 + \frac{3n(n+1)(2n+1)}{6} + \frac{2n(n+1)}{2} \\
&= \left[\frac{n(n+1)}{2} \right]^2 + \frac{n(n+1)(2n+1)}{2} + n(n+1) \\
&= \frac{n(n+1)}{2} \left[\frac{n(n+1)}{2} + 2n+1+2 \right] \\
&= \frac{n(n+1)}{2} \left[\frac{n^2 + n + 4n + 6}{2} \right] \\
&= \frac{n(n+1)}{4} (n^2 + 5n + 6) \\
&= \frac{n(n+1)}{4} (n^2 + 2n + 3n + 6) \\
&= \frac{n(n+1)[n(n+2) + 3(n+2)]}{4} \\
&= \frac{n(n+1)(n+2)(n+3)}{4}
\end{aligned}$$

Page : 196 , Block Name : Exercise 9.4

Q3 Find the sum to n terms of the series $3 \times 1^2 + 5 \times 2^2 + 7 \times 3^2 + \dots$

Answer.

The given series is $3 \times 1^2 + 5 \times 2^2 + 7 \times 3^2 + \dots$

n^{th} term, $a_n = (2n+1)n^2 = 2n^3 + n^2$

$$\begin{aligned}
\therefore S_n &= \sum_{k=1}^n a_k \\
&= \sum_{k=1}^n (2k^3 + k^2) = 2 \sum_{k=1}^n k^3 + \sum_{k=1}^n k^2 \\
&= 2 \left[\frac{n(n+1)}{2} \right]^2 + \frac{n(n+1)(2n+1)}{6} \\
&= \frac{n^2(n+1)^2}{2} + \frac{n(n+1)(2n+1)}{6} \\
&= \frac{n(n+1)}{2} \left[n(n+1) + \frac{2n+1}{3} \right]
\end{aligned}$$

$$\begin{aligned}
&= \frac{n(n+1)}{2} \left[\frac{3n^2 + 3n + 2n + 1}{3} \right] \\
&= \frac{n(n+1)}{2} \left[\frac{3n^2 + 5n + 1}{3} \right] \\
&= \frac{n(n+1)(3n^2 + 5n + 1)}{6}
\end{aligned}$$

Page : 196 , Block Name : Exercise 9.4

Q4 Find the sum to n terms of the series $\frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots$

Answer, The given series is $\frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots$

n^{th} term, $a_n = \frac{1}{n(n+1)}$

$$= \frac{1}{n} - \frac{1}{n+1} \quad [\text{by partial fraction}]$$

$$a_1 = \frac{1}{1} - \frac{1}{2}$$

$$a_2 = \frac{1}{2} - \frac{1}{3}$$

$$a_3 = \frac{1}{3} - \frac{1}{4}$$

$$a_3 = \frac{1}{3} - \frac{1}{4} \dots$$

$$a_n = \frac{1}{3} - \frac{1}{4+1}$$

Adding the above terms column wise, we obtain

$$a_1 + a_2 + \dots + a_n = \left[\frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} \right] - \left[\frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n+1} \right]$$

$$\therefore S_n = 1 - \frac{1}{n+1} = \frac{n+1-1}{n+1} = \frac{n}{n+1}$$

Page : 196 , Block Name : Exercise 9.4

Q5 Find the sum to n terms of the series $5^2 + 6^2 + 7^2 + \dots + 20^2$

Answer.

The given series is $5^2 + 6^2 + 7^2 + \dots + 20^2$

n^{th} term, $a_n = (n+4)^2 = n^2 + 8n + 16$

$$\begin{aligned}
\therefore S_n &= \sum_{k=1}^n a_k = \sum_{k=1}^n (k^2 + 8k + 16) \\
&= \sum_{k=1}^n k^2 + 8 \sum_{k=1}^n k + \sum_{k=1}^n 16 \\
&= \frac{n(n+1)(2n+1)}{6} + \frac{8n(n+1)}{2} + 16n
\end{aligned}$$

$$\begin{aligned}
& 16^{\text{th}} \text{ term is } (16 + 4)^2 = 20^2 \\
\therefore S_{16} &= \frac{16(16+1)(2 \times 16+1)}{6} + \frac{8 \times 16 \times (16+1)}{2} + 16 \times 16 \\
&= \frac{(16)(17)(33)}{6} + \frac{(8) \times 16 \times (16+1)}{2} + 16 \times 16 \\
&= \frac{(16)(17)(33)}{6} + \frac{(8)(16)(17)}{2} + 256 \\
&\quad = 1496 + 1088 + 256 \\
&\quad = 2840 \\
\therefore 5^2 + 6^2 + 7^2 + \dots + 20^2 &= 2840
\end{aligned}$$

Page : 196 , Block Name : Exercise 9.4

Q6 Find the sum to n terms of the series $3 \times 8 + 6 \times 11 + 9 \times 14 + \dots$

Answer. The given series is $3 \times 8 + 6 \times 11 + 9 \times 14 + \dots$

$$\begin{aligned}
a_n &= (n^{\text{th}} \text{ term of } 3, 6, 9 \dots) \times (n^{\text{th}} \text{ term of } 8, 11, 14, \dots) \\
&= (3n)(3n + 5) \\
&= 9n^2 + 15n \\
\therefore S_n &= \sum_{k=1}^n a_k = \sum_{k=1}^n (9k^2 + 15k) \\
&= 9 \sum_{k=1}^n k^2 + 15 \sum_{k=1}^n k \\
&= 9 \times \frac{n(n+1)(2n+1)}{6} + 15 \times \frac{n(n+1)}{2} \\
&= \frac{3n(n+1)(2n+1)}{2} + \frac{15n(n+1)}{2} \\
&= \frac{3n(n+1)}{2} (2n+1+5) \\
&= \frac{3n(n+1)}{2} (2n+6) \\
&= 3n(n+1)(n+3)
\end{aligned}$$

Page : 196 , Block Name : Exercise 9.4

Q7 Find the sum to n terms of the series $1^2 + (1^2 + 2^2) + (1^2 + 2^2 + 3^2) + \dots$

Answer.

The given series $|s1^2 + (1^2 + 2^2) + (1^2 + 2^2 + 3^2) + \dots$

$$\begin{aligned}
 a_n &= (1^2 + 2^2 + 3^2 + \dots + n^2) \\
 &= \frac{n(n+1)(2n+1)}{6} \\
 &= \frac{n(2n^2+3n+1)}{6} = \frac{2^3+3n^2+n}{6} \\
 &= \frac{1}{3}n^3 + \frac{1}{2}n^2 + \frac{1}{6}n \\
 \therefore S_n &= \sum_{k=1}^n a_k \\
 &= \sum_{k=1}^n \left(\frac{1}{3}k^3 + \frac{1}{2}k^2 + \frac{1}{6}k \right) \\
 &= \frac{1}{3} \sum_{k=1}^n k^3 + \frac{1}{2} \sum_{k=1}^n k^2 + \frac{1}{6} \sum_{k=1}^n k \\
 &= \frac{1}{3} \frac{n^2(n+1)^2}{(2)^2} + \frac{1}{2} \times \frac{n(n+1)(2n+1)}{6} + \frac{1}{6} \times \frac{n(n+1)}{2} \\
 &= \frac{n(n+1)}{6} \left[\frac{n(n+1)}{2} + \frac{(2n+1)}{2} + \frac{1}{2} \right] \\
 &= \frac{n(n+1)}{6} \left[\frac{n^2+n+2n+1+1}{2} \right] \\
 &= \frac{n(n+1)}{6} \left[\frac{n^2+n+2n+2}{2} \right] \\
 &= \frac{n(n+1)}{6} \left[\frac{n(n+1)+2(n+1)}{2} \right] \\
 &= \frac{n(n+1)}{6} \left[\frac{(n+1)(n+2)}{2} \right] \\
 &= \frac{n(n+1)^2(n+1)(n+2)}{2} \\
 &= \frac{n(n+1)^2(n+1)(n+2)}{12}
 \end{aligned}$$

Page : 196 , Block Name : Exercise 9.4

Q8 Find the sum to n terms of the series $n(n+1)(n+4)$

Answer.

$$\begin{aligned}
 a_n &= n(n+1)(n+4) = n(n^2+5n+4) = n^3+5n^2+4n \\
 \therefore S_n &= \sum_{k=1}^n a_k = \sum_{k=1}^n k^3 + 5 \sum_{k=1}^n k^2 + 4 \sum_{i=1}^n k \\
 &= \frac{n^2(n+1)^2}{4} + \frac{5n(n+1)(2n+1)}{6} + \frac{4n(n+1)}{2}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{n(n+1)}{2} \left[\frac{n(n+1)}{2} + \frac{5(2n+1)}{3} + 4 \right] \\
&= \frac{n(n+1)}{2} \left[\frac{3n^2 + 3n + 20n + 10 + 24}{6} \right] \\
&= \frac{n(n+1)}{2} \left[\frac{3n^2 + 23n + 34}{6} \right] \\
&= \frac{n(n+1)(3n^2 + 23n + 34)}{12}
\end{aligned}$$

Page : 196 , Block Name : Exercise 9.4

Q9 Find the sum to n terms of the series $n^2 + 2^n$

Answer.

$$a_n = n^2 + 2^n$$

$$\therefore S_n = \sum_{k=1}^n k^2 + 2^k = \sum_{k=1}^n k^2 + \sum_{k=1}^n 2^k \quad \dots (1)$$

$$\text{Consider } \sum_{k=1}^n 2^k = 2^1 + 2^2 + 2^3 + \dots$$

The above series $2, 2^2, 2^3, \dots$ is a G.P. with both the first term and common ratio equal to 2 .

$$\therefore \sum_{k=1}^n 2^k = \frac{(2)[(2)^n - 1]}{2 - 1} = 2(2^n - 1) \quad \dots (2)$$

Therefore, from (1) and (2), we obtain

$$S_n = \sum_{k=1}^n k^2 + 2(2^n - 1) = \frac{n(n+1)(2n+1)}{6} + 2(2^n - 1)$$

Page : 196 , Block Name : Exercise 9.4

Q10 Find the sum to n terms of the series $(2n - 1)^2$

Answer.

$$a_n = (2n - 1)^2 = 4n^2 - 4n + 1$$

$$\therefore S_n = \sum_{k=1}^n a_k = \sum_{k=1}^n (4k^2 - 4k + 1)$$

$$= 4 \sum_{k=1}^n k^2 - 4 \sum_{k=1}^n k + \sum_{k=1}^n 1$$

$$= \frac{4n(n+1)(2n+1)}{6} - \frac{4n(n+1)}{2} + n$$

$$\begin{aligned}
&= \frac{2n(n+1)(2n+1)}{3} - 2n(n+1) + n \\
&= n \left[\frac{2(2n^2 + 3n + 1)}{3} - 2(n+1) + 1 \right] \\
&= n \left[\frac{4n^2 + 6n + 2 - 6n - 6 + 3}{3} \right] \\
&= n \left[\frac{4n^2 - 1}{3} \right] \\
&= \frac{n(2n+1)(2n-1)}{3} \\
&= \frac{n(2n+1)(2n-1)}{3}
\end{aligned}$$

Page : 196 , Block Name : Exercise 9.4

Miscellaneous Exercise

Q1 Show that the sum of $(m+n)^{th}$ and $(m-n)^{th}$ terms of an A.P. is equal to twice the m^{th} term.

Answer. Let a and d be the first term and the common difference of the A.P. respectively.

It is known that the k th term of an A. P. is given by

$$a_k = a + (k-1)d$$

$$\therefore a_{m+n} = a + (m+n-1)d$$

$$a_{m-n} = a + (m-n-1)d$$

$$a_m = a + (m-1)d$$

$$\therefore a_{m+n} + a_{m-n} = a + (m+n-1)d + a + (m-n-1)d$$

$$= 2a + (m+n-1 + m-n-1)d$$

$$= 2a + (2m-2)d$$

$$= 2a + 2(m-1)d$$

$$= 2[a + (m-1)d]$$

$$= 2a_m$$

Thus, the sum of $(m+n)^{th}$ and $(m-n)^{th}$ terms of an A.P. is equal to twice the m^{th} term.

Page : 199 , Block Name : Miscellaneous Exercise

Q2 If the sum of three numbers in A.P., is 24 and their product is 440, find the numbers.

Answer. Let the three numbers in A.P. be $a-d$, a , and $a+d$.

According to the given information,

$$(a-d) + (a) + (a+d) = 24 \dots (1)$$

$$\Rightarrow 3a = 24$$

$$\therefore a = 8$$

$$(a - d) a (a + d) = 440 \dots (2)$$

$$\Rightarrow (8 - d) (8) (8 + d) = 440$$

$$\Rightarrow (8 - d) (8 + d) = 55$$

$$\Rightarrow 64 - d^2 = 55$$

$$\Rightarrow d^2 = 64 - 55 = 9$$

$$\Rightarrow d = \pm 3$$

Therefore, when $d = 3$, the numbers are 5, 8, and 11 and when $d = -3$, the numbers are 11, 8, and 5.

Thus, the three numbers are 5, 8, and 11.

Page : 199 , Block Name : Miscellaneous Exercise

Q3 Let the sum of $n, 2n, 3n$ terms of an A.P. be S_1, S_2, S_3 , respectively, show that $S_3 = 3(S_2 - S_1)$

Answer. Let a and b be the first term and the common difference of the A.P. respectively.

Therefore,

$$S_1 = \frac{n}{2} [2a + (n - 1)d] \dots (1)$$

$$S_2 = \frac{2n}{2} [2a + (2n - 1)d] = n[2a + (2n - 1)d] \dots (2)$$

$$S_3 = \frac{3n}{2} [2a + (3n - 1)d] \dots (3)$$

From (1) and (2), we obtain

$$S_2 - S_1 = n[2a + (2n - 1)d] - \frac{n}{2} [2a + (n - 1)d]$$

$$= n \left\{ \frac{4a + 4nd - 2d - 2a - nd + d}{2} \right\}$$

$$= n \left[\frac{2a + 3nd - d}{2} \right]$$

$$= \frac{n}{2} [2a + (3n - 1)d]$$

$$\therefore 3(S_2 - S_1) = \frac{3n}{2} [2a + (3n - 1)d] = S_3 \quad [\text{From (3)}]$$

Hence, the given result is proved.

Page : 199 , Block Name : Miscellaneous Exercise

Q4 Find the sum of all numbers between 200 and 400 which are divisible by 7.

Answer. The numbers lying between 200 and 400, which are divisible by 7, are

203, 210, 217, ... 399

\therefore First term, $a = 203$

Last term, $l = 399$

Common difference, $d = 7$

Let the number of terms of the A.P. be n .

$$\therefore an = 399 = a + (n - 1)d$$

$$\Rightarrow 399 = 203 + (n - 1)7$$

$$\Rightarrow 7(n - 1) = 196$$

$$\Rightarrow n - 1 = 28$$

$$\Rightarrow n = 29$$

$$\begin{aligned}\therefore S_{29} &= \frac{29}{2}(203 + 399) \\ &= \frac{29}{2}(602) \\ &= (29)(301) \\ &= 8729\end{aligned}$$

Thus, the required sum is 8729

Page : 199 , Block Name : Miscellaneous Exercise

Q5 Find the sum of integers from 1 to 100 that are divisible by 2 or 5.

Answer. The integers from 1 to 100, which are divisible by 2, are 2, 4, 6... 100. This forms an A.P. with both the first term and common difference equal to 2.

$$\Rightarrow 100 = 2 + (n - 1) 2$$

$$\Rightarrow n = 50$$

$$\begin{aligned}\therefore 2 + 4 + 6 + \dots + 100 &= \frac{50}{2}[2(2) + (50 - 1)(2)] \\ &= \frac{50}{2}[4 + 98] \\ &= (25)(102) \\ &= 2550\end{aligned}$$

The integers from 1 to 100, which are divisible by 5, are 5, 10... 100.

This forms an A.P. with both the first term and common difference equal to 5.

$$\therefore 100 = 5 + (n - 1) 5$$

$$\Rightarrow 5n = 100$$

$$\Rightarrow n = 20$$

$$\begin{aligned}\therefore 5 + 10 + \dots + 100 &= \frac{20}{2}[2(5) + (20 - 1)5] \\ &= 10[10 + (19)5] \\ &= 10[10 + 95] = 10 \times 105 \\ &= 1050\end{aligned}$$

The integers, which are divisible by both 2 and 5, are 10, 20, ... 100.

This also forms an A.P. with both the first term and common difference equal to 10.

$$\therefore 100 = 10 + (n - 1) (10)$$

$$\Rightarrow 100 = 10n$$

$$\Rightarrow n = 10$$

$$\begin{aligned}\therefore 10 + 20 + \dots + 100 &= \frac{10}{2}[2(10) + (10 - 1)(10)] \\ &= 5[20 + 90] = 5(110) = 550\end{aligned}$$

$$\therefore \text{Required sum} = 2550 + 1050 - 550 = 3050$$

Thus, the sum of the integers from 1 to 100, which are divisible by 2 or 5, is 3050.

Page : 199 , Block Name : Miscellaneous Exercise

Q6 Find the sum of all two digit numbers which when divided by 4, yields 1 as Remainder.

Answer. The two-digit numbers, which when divided by 4, yield 1 as remainder, are 13, 17, ... 97.

This series forms an A.P. with first term 13 and common difference 4.

Let n be the number of terms of the A.P.

It is known that the n th term of an A.P. is given by, $a_n = a + (n - 1) d$

$$\therefore 97 = 13 + (n - 1) (4)$$

$$\Rightarrow 4 (n - 1) = 84$$

$$\Rightarrow n - 1 = 21$$

$$\Rightarrow n = 22$$

Sum of n terms of an A.P. is given by,

$$S_n = \frac{n}{2} [2a + (n - 1)d]$$

$$\begin{aligned} \therefore S_{22} &= \frac{22}{2} [22(13) + (22 - 1)(4)] \\ &= 11[26 + 84] \\ &= 1210 \end{aligned}$$

Thus, the required sum is 1210.

Page : 199 , Block Name : Miscellaneous Exercise

Q7 If f is a function satisfying $f(x + y) = f(x) f(y)$ for all $x, y \in \mathbb{N}$ such that $f(1) = 3$ and $\sum_{x=1}^n f(x) = 120$, find the value of n

Answer. It is given that,

$$f(x + y) = f(x) \times f(y) \text{ for all } x, y \in \mathbb{N} \dots (1)$$

$$f(1) = 3$$

Taking $x = y = 1$ in (1), we obtain

$$f(1 + 1) = f(2) = f(1) f(1) = 3 \times 3 = 9$$

Similarly,

$$f(1 + 1 + 1) = f(3) = f(1 + 2) = f(1) f(2) = 3 \times 9 = 27$$

$$f(4) = f(1 + 3) = f(1) f(3) = 3 \times 27 = 81$$

$\therefore f(1), f(2), f(3), \dots$, that is 3, 9, 27, ..., forms a G.P. with both the first term and common ratio equal to 3.

$$\text{It is known that, } S_n = \frac{a(r^n - 1)}{r - 1}$$

It is given that, $\sum_{x=1}^n f(x) = 120$

$$\therefore 120 = \frac{3(3^n - 1)}{3 - 1}$$

$$\Rightarrow 120 = \frac{3}{2} (3^n - 1)$$

$$\Rightarrow 3^n - 1 = 80$$

$$\Rightarrow 3^n = 81 = 3^4$$

$$\therefore n = 4$$

Thus, the value of n is 4.

Page : 199 , Block Name : Miscellaneous Exercise

Q8 The sum of some terms of G.P. is 315 whose first term and the common ratio are 5 and 2,

respectively.

Find the last term and the number of terms.

Answer. Let the sum of n terms of the G.P. be 315.

It is known that, $S_n = \frac{a(r^n-1)}{r-1}$

It is given that the first term a is 5 and common ratio r is 2.

$$\therefore 315 = \frac{5(2^n-1)}{2-1}$$

$$\Rightarrow 2^n - 1 = 63$$

$$\Rightarrow 2^n = 64 = (2)^6$$

$$\Rightarrow n = 6$$

\therefore Last term of the G.P = 6th term = $ar^{n-1} = (5)(2)^5 = (5)(32) = 160$

\therefore Last term of the G.P = 6th term = $ar^{n-1} = (5)(2)^5 = (5)(32) = 160$

Thus, the last term of the G.P. is 160.

Page : 199 , Block Name : Miscellaneous Exercise

Q9 The first term of a G.P. is 1. The sum of the third term and fifth term is 90.

Find the common ratio of G.P.

Answer. Let a and r be the first term and the common ratio of the G.P. respectively.

$$\therefore a = 1$$

$$a_3 = ar^2 = r^2$$

$$a_5 = ar^4 = r^4$$

$$\therefore r^2 + r^4 = 90$$

$$\Rightarrow r^4 + r^2 - 90 = 0$$

$$\Rightarrow r^2 = \frac{-1 \pm \sqrt{1+360}}{2} = \frac{-1 \pm \sqrt{361}}{2} = \frac{-1 \pm 19}{2} = -10 \text{ or } 9$$

$$\therefore r = \pm 3 \quad (\text{Taking real roots})$$

Thus, the common ratio of the G.P. is ± 3 .

Page : 199 , Block Name : Miscellaneous Exercise

Q10 The sum of three numbers in G.P. is 56. If we subtract 1, 7, 21 from these numbers in that order, we obtain an arithmetic progression. Find the numbers.

Answer.

Let the three numbers in G.P. be a , ar , and ar^2 .

From the given condition, $a + ar + ar^2 = 56$

$$\Rightarrow a(1 + r + r^2) = 56$$

$$\Rightarrow a = \frac{56}{1+r+r^2} \dots (1)$$

$a - 1$, $ar - 7$, $ar^2 - 21$ forms an A.P.

$$\therefore (ar - 7) - (a - 1) = (ar^2 - 21) - (ar - 7)$$

$$\Rightarrow ar - a - 6 = ar^2 - ar - 14$$

$$\Rightarrow ar^2 - 2ar + a = 8$$

$$\Rightarrow ar^2 - ar - ar + a = 8$$

$$\Rightarrow a(r - 1)^2 = 8 \dots (2)$$

$$\Rightarrow \frac{56}{1+r+r^2}(r - 1)^2 = 8$$

$$\Rightarrow 7 + r + r^2$$

$$\Rightarrow 7r^2 - 14r + 7 - 1 + r + r^2$$

$$\Rightarrow 7r^2 - 14r + 7 - 1 - r - r^2 = 0$$

$$\Rightarrow 6r^2 - 15r + 6 = 0$$

$$\Rightarrow 6r^2 - 12r - 3r + 6 = 0$$

$$\Rightarrow 6r(r - 2) - 3(r - 2) = 0$$

$$\Rightarrow (6r - 3)(r - 2) = 0$$

$$\therefore r = 2, \frac{1}{2}$$

When $r = 2, a = 8$

When $r = \frac{1}{2}, a = 32$

Therefore, when $r = 2$, the three numbers in G.P. are 8, 16, and 32.

When $r = \frac{1}{2}$, the three numbers in G.P. are 32, 16, and 8.

Thus, in either case, the three required numbers are 8, 16, and 32.

Page : 199 , Block Name : Miscellaneous Exercise

Q11 A G.P. consists of an even number of terms.

If the sum of all the terms is 5 times the sum of terms occupying odd places, then find its common ratio.

Answer.

Let the G.P. be $T_1, T_2, T_3, \dots, T_4, \dots, T_{2n}$.

Number of terms = $2n$

According to the given condition,

$$T_1 + T_2 + T_3 + \dots + T_{2n} = 5[T_1 + T_3 + \dots + T_{2n-1}] = 0$$

$$\Rightarrow T_1 + T_2 + T_3 + \dots + T_{2n} - 5[T_1 + T_3 + \dots + T_{2n-1}] = 0$$

$$\Rightarrow T_2 + T_4 + \dots + T_{2n} = 4[T_1 + T_3 + \dots + T_{2n-1}] = 0$$

Let the G.P. be a, ar, ar^2, ar^3, \dots

$$\therefore \frac{ar(r^n - 1)}{r - 1} = \frac{4 \times a(r^n - 1)}{r - 1}$$

$$\Rightarrow ar = 4a$$

$$\Rightarrow r = 4$$

Thus, the common ratio of the G.P. is 4.

Page : 199 , Block Name : Miscellaneous Exercise

Q12 The sum of the first four terms of an A.P. is 56. The sum of the last four terms is 112.

If its first term is 11, then find the number of terms.

Answer. Let the A.P. be $a, a + d, a + 2d, a + 3d, \dots, a + (n - 2)d, a + (n - 1)d$.

Sum of first four terms $= a + (a + d) + (a + 2d) + (a + 3d) = 4a + 6d$

Sum of last four terms $= [a + (n - 4)d] + [a + (n - 3)d] + [a + (n - 2)d] + [a + (n - 1)d]$

$= 4a + (4n - 10)d$

According to the given condition,

$$4a + 6d = 56$$

$$\Rightarrow 4(11) + 6d = 56 \text{ [Since } a = 11 \text{ (given)]}$$

$$\Rightarrow 6d = 12$$

$$\Rightarrow d = 2$$

$$\therefore 4a + (4n - 10)d = 112$$

$$\Rightarrow 4(11) + (4n - 10)2 = 112$$

$$\Rightarrow (4n - 10)2 = 68$$

$$\Rightarrow 4n - 10 = 34$$

$$\Rightarrow 4n = 44$$

$$\Rightarrow n = 11$$

Thus, the number of terms of the A.P. is 11.

Page : 199 , Block Name : Miscellaneous Exercise

Q13 If $\frac{a+bx}{a-bx} = \frac{b+cx}{b-cx} = \frac{c+dx}{c-dx} (x \neq 0)$ then show that a, b, c and d are in G.P.

Answer.

It is given that,

$$\frac{a+bx}{a-bx} = \frac{b+cx}{b-cx}$$

$$\Rightarrow (a + bx)(b - cx) = (b + cx)(a - bx)$$

$$\Rightarrow ab - acx + b^2x - bcx^2 = ab - b^2x + acx - bcx^2$$

$$\Rightarrow b^2x = 2acx$$

$$\Rightarrow b^2 = ac$$

$$\Rightarrow \frac{b}{a} = \frac{c}{b} \quad \dots (1)$$

$$\text{Also, } \frac{b+cx}{b-cx} = \frac{c+dx}{c-dx}$$

$$\Rightarrow (b + cx)(c - dx) = (b - cx)(c + dx)$$

$$\Rightarrow bc - bdx + c^2x - cdx^2 = bc + bdx - c^2x - cdx^2$$

$$\Rightarrow 2c^2x = 2bdx$$

$$\Rightarrow c^2 = bd$$

$$\Rightarrow \frac{c}{d} = \frac{d}{c} \quad \dots (2)$$

From (1) and (2), we obtain

$$\frac{b}{a} = \frac{c}{b} = \frac{d}{c}$$

Thus, a, b, c , and d are in G.P.

Page : 199 , Block Name : Miscellaneous Exercise

Q14 Let S be the sum, P the product and R the sum of reciprocals of n terms in a G.P.
Prove that $P^2 R^n = S^n$

Answer.

Let the G.P. be $a, ar, ar^2, ar^3, \dots, ar^{n-1} \dots$

According to the given information,

$$\begin{aligned}
 S &= \frac{a(r^n - 1)}{r - 1} \\
 P &= a^n \times r^{1+2+\dots+n-1} \\
 &= a^n r^{\frac{n-1}{2}} \left[\because \text{Sum of first } n \text{ natural numbers is } n \frac{(n+1)}{2} \right] \\
 R &= \frac{1}{a} + \frac{1}{ar} + \dots + \frac{1}{ar^{n-1}} \\
 &= \frac{r^{n-1} + r^{n-2} + \dots + r + 1}{ar^{n-1}} \\
 &= \frac{1(r^n - 1)}{(r - 1)} \times \frac{1}{ar^{n-1}} \quad [\because 1, r, \dots, r^{n-1} \text{ from a G.P.}] \\
 &= \frac{r^n - 1}{ar^{n-1}(r - 1)} \\
 \therefore P^2 R^n &= a^{2n} r^{n(n-1)} \frac{(r^n - 1)^n}{a^n r^{n(n-1)} (r - 1)^n} \\
 &= \frac{a^n (r^n - 1)^n}{(r - 1)^n} \\
 &= \left[\frac{a(r^n - 1)}{(r - 1)} \right]^n \\
 &= S^n
 \end{aligned}$$

Hence, $P^2 R^n = S^n$

Page : 199 , Block Name : Miscellaneous Exercise

Q15 The p^{th} , q^{th} and r^{th} terms of an A.P. are a, b, c, respectively. Show that $(q - r)a + (r - p)b + (p - q)c = 0$

Answer. Let t and d be the first term and the common difference of the A.P. respectively.

The nth term of an A.P. is given by, $a_n = t + (n - 1)d$

Therefore,

$$ap = t + (p - 1)d = a \dots (1)$$

$$aq = t + (q - 1)d = b \dots (2)$$

$$ar = t + (r - 1)d = c \dots (3)$$

Subtracting equation (2) from (1), we obtain

$$(p - 1 - q + 1)d = a - b$$

$$\Rightarrow (p - q)d = a - b$$

$$\therefore d = \frac{a - b}{p - q} \dots (4)$$

Subtracting equation (3) from (2), we obtain

$$(q - 1 - r + 1)d = b - c$$

$$\Rightarrow (q - r)d = b - c$$

$$\Rightarrow d = \frac{b-c}{q-r} \dots (5)$$

Equating both the values of d obtained in (4) and (5), we obtain

$$\frac{a-b}{p-q} = \frac{b-c}{q-r}$$

$$\Rightarrow (a-b)(q-r) = (b-c)(p-q)$$

$$\Rightarrow aq - bq - ar + br = bp - cp + cq$$

$$\Rightarrow bp - cp + cq - aq + ar - br = 0$$

$$\Rightarrow (-aq + ar) + (bp - br) + (-cp + cq) = 0$$

$$\Rightarrow -a(q-r) - b(r-p) - c(p-q) = 0$$

$$\Rightarrow a(q-r) + b(r-p) + c(p-q) = 0$$

Thus, the given result is proved.

Page : 199 , Block Name : Miscellaneous Exercise

Q16 If $a\left(\frac{1}{b} + \frac{1}{c}\right), b\left(\frac{1}{c} + \frac{1}{a}\right), c\left(\frac{1}{a} + \frac{1}{b}\right)$ are in A.P., prove that a, b, c are in A.P.

Answer.

It is given that $a\left(\frac{1}{b} + \frac{1}{c}\right), b\left(\frac{1}{c} + \frac{1}{a}\right), c\left(\frac{1}{a} + \frac{1}{b}\right)$ are in A.P.

$$\therefore b\left(\frac{1}{c} + \frac{1}{a}\right) - a\left(\frac{1}{b} + \frac{1}{c}\right) = c\left(\frac{1}{a} + \frac{1}{b}\right) - b\left(\frac{1}{c} + \frac{1}{a}\right)$$

$$\Rightarrow \frac{b(a+c)}{ac} - \frac{a(b+c)}{bc} = \frac{c(a+b)}{ab} - \frac{b(a+c)}{ac}$$

$$\Rightarrow \frac{b^2a + b^2c - a^2b - a^2c}{abc} = \frac{c^2a + c^2b - b^2a - b^2c}{abc}$$

$$\Rightarrow b^2a - a^2b + b^2c - a^2c = c^2a - b^2a + c^2b - b^2c$$

$$\Rightarrow ab(b-a) + c(b^2 - a^2) = a(c^2 - b^2) + bc(c-b)$$

$$\Rightarrow (b-a)(ab + cb + ca) = (c-b)(c+b) + bc(c-b)$$

$$\Rightarrow (b-a)(ab + c - b) = (c-b)(ac + ab + bc)$$

$$\Rightarrow b - a = c - b$$

Thus, a, b, and c are in A.P.

Page : 199 , Block Name : Miscellaneous Exercise

Q17 If a, b, c, d are in G.P, prove that $(a^n + b^n), (b^n + c^n), (c^n + d^n)$ are in G.P.

Answer. It is given that a, b, c, and d are in G.P.

$$\therefore b^2 = ac \dots (1)$$

$$c^2 = bd \dots (2)$$

$$ad = bc \dots (3)$$

It has to be proved that $(a^n + b^n), (b^n + c^n), (c^n + d^n)$ are in G.P. i.e.,

$$(b^n + c^n)^2 = (a^n + b^n)(c^n + d^n)$$

Consider L.H.S.

$$(b^n + c^n)^2 = b^{2n} + 2b^n c^n + c^{2n}$$

$$= (b^2)^n + 2b^n c^n + (c^2)^n$$

$$= (ac)^n + 2b^n c^n + (bd)^n [U \text{ sing(1) and (2)}]$$

$$= a^n c^n + b^n c^n + b^n c^n + b^n d^n$$

$$= a^n c^n + b^n c^n + a^n d^n + b^n d^n [\text{Using (3)}]$$

$$= c^n (a^n + b^n) + d^n (a^n + b^n)$$

$$= (a^n + b^n)(c^n + d^n)$$

= R. H. S.

$$\therefore (b^n + c^n)^2 = (a^n + b^n)(c^n + d^n)$$

Thus, $(a^n + b^n)$, $(b^n + c^n)$, and $(c^n + d^n)$ are in G.P.

Page : 199 , Block Name : Miscellaneous Exercise

Q18

If a and b are the roots of $x^2 - 3x + p = 0$ and c, d are roots of $x^2 - 12x + q = 0$ where a, b, c, d form a G.P. Prove that $(q + p) : (q - p) = 17 : 15$.

Answer.

It is given that a and b are the roots of $x^2 - 3x + p = 0$

$$\therefore a + b = 3 \text{ and } ab = p \dots (1)$$

Also, c and d are the roots of $x^2 - 12x + q = 0$

$$\therefore c + d = 12 \text{ and } cd = q \dots (2)$$

It is given that a, b, c, d are in G.P.

$$\text{Let } a = x, b = xr, c = xr^2, d = xr^3$$

From (1) and (2), we obtain

$$x + xr = 3$$

$$\Rightarrow x(1 + r) = 3$$

$$xr^2 + xr^3 = 12$$

$$\Rightarrow x r^2(1 + r) = 12$$

On dividing, we obtain

$$\frac{xr^2(1+r)}{x(1+r)} = \frac{12}{3}$$

$$\Rightarrow r^2 = 4$$

$$\Rightarrow r^2 = 4$$

$$\Rightarrow r = \pm 2$$

$$\text{When } r = -2, x = \frac{3}{1-2} = \frac{3}{-1} = -3$$

$$\text{When } r = 2, x = \frac{3}{1-2} = \frac{3}{-1} = -3$$

Case : I

When $r = 2$ and $x = 1$

$$ab = x^2r = 2$$

$$cd = x^2r^5 = 32$$

$$\therefore \frac{q+p}{q-p} = \frac{32+2}{32-2} = \frac{34}{30} = \frac{17}{15}$$

$$\text{i.e., } (q+p) : (q-p) = 17 : 15$$

Case : II

When $r = -2$, $x = -3$

$$ab = x^2r = -18$$

$$cd = x^2r^5 = -288$$

$$\therefore \frac{q+p}{q-p} = \frac{-288-18}{-288+18} = \frac{-306}{-270} = \frac{17}{15}$$

$$\text{i.e., } (q+p) : (q-p) = 17 : 15$$

Thus, in both the cases, we obtain $(q+p) : (q-p) = 17 : 15$

Page : 199 , Block Name : Miscellaneous Exercise

Q19 The ratio of the A.M. and G.M. of two positive numbers a and b , is $m : n$.

Show that $a : b = \left(m + \sqrt{m^2 - n^2}\right) : \left(m - \sqrt{m^2 - n^2}\right)$

Answer. Let the two numbers be a and b .

$$\text{A.M.} = \frac{a+b}{2} \text{ and G.M.} = \sqrt{ab}$$

According to the given condition,

$$\frac{a+b}{2\sqrt{ab}} = \frac{m}{n}$$

$$\Rightarrow \frac{(a+b)^2}{4(ab)} = \frac{m^2}{n^2}$$

$$\Rightarrow (a+b)^2 = \frac{4abm^2}{n^2}$$

$$\Rightarrow (a+b) = \frac{2\sqrt{ab}m}{n}$$

Using this in the identity $(a-b)^2 = (a+b)^2 - 4ab$, we obtain

$$(a-b)^2 = \frac{4abm^2}{n^2} - 4ab = \frac{4ab(m^2-n^2)}{n^2}$$

$$\Rightarrow (a-b) = \frac{2\sqrt{ab}\sqrt{m^2-n^2}}{n} \quad \dots (2)$$

Adding (1) and (2), we obtain

$$2a = \frac{2\sqrt{ab}}{n} \left(m + \sqrt{m^2 - n^2}\right)$$

$$\Rightarrow a = \frac{\sqrt{ab}}{n} \left(m + \sqrt{m^2 - n^2}\right)$$

Substituting the value of a in (1), we obtain

$$b = \frac{2\sqrt{ab}}{n}m - \frac{\sqrt{ab}}{n} \left(m + \sqrt{m^2 - n^2}\right)$$

$$\begin{aligned}
&= \frac{\sqrt{ab}}{n} m - \frac{\sqrt{ab}}{n} \sqrt{m^2 - n^2} \\
&= \frac{\sqrt{ab}}{n} (m - \sqrt{m^2 - n^2}) \\
\therefore a : b &= \frac{a}{b} = \frac{\frac{\sqrt{ab}}{\sqrt{ab}} (m + \sqrt{m^2 - n^2})}{n} = \frac{(m + \sqrt{m^2 - n^2})}{(m - \sqrt{m^2 - n^2})} \\
\text{Thus, } a : b &= (m + \sqrt{m^2 - n^2}) : (m - \sqrt{m^2 - n^2})
\end{aligned}$$

Page : 200 , Block Name : Miscellaneous Exercise

Q20 If a, b, c are in A.P.; b, c, d are in G.P. and $\frac{1}{c}, \frac{1}{d}, \frac{1}{e}$ are in A.P. prove that a, c, e are in G.P.

Answer. It is given that a, b, c are in A.P.

$$\therefore b - a = c - b \dots (1)$$

It is given that b, c, d, are in G.P.

$$\therefore c^2 = bd \dots (2)$$

Also, $\frac{1}{c}, \frac{1}{d}, \frac{1}{e}$ are in A.P.

$$\frac{1}{d} - \frac{1}{c} = \frac{1}{e} - \frac{1}{d}$$

$$\frac{2}{d} = \frac{1}{c} + \frac{1}{e} \dots (3)$$

It has to be proved that a, c, e are in G.P. i.e., $a^2 = ae$

From (1), we obtain

$$2b = a + c$$

$$\Rightarrow b = \frac{a+c}{2}$$

From (2), we obtain

$$d = \frac{c^2}{b}$$

Substituting these values in (3), we obtain

$$\frac{2b}{c^2} = \frac{1}{c} + \frac{1}{e}$$

$$\Rightarrow \frac{2(a+c)}{2c^2} = \frac{1}{c} + \frac{1}{e}$$

$$\Rightarrow \frac{a+c}{c^2} = \frac{e+c}{ce}$$

$$\Rightarrow \frac{a+c}{c} = \frac{e+c}{c}$$

$$\Rightarrow (a+c)e = (e+c)c$$

$$\Rightarrow ae + ce = ec + c^2$$

$$\Rightarrow c^2 = ae$$

Thus, a, c, and e are in G.P.

Page : 200 , Block Name : Miscellaneous Exercise

Q21 Find the sum of the following series up to n terms:

(i) $5 + 55 + 555 + \dots$ (ii) $.6 + .66 + .666 + \dots$

Answer. (i) $5 + 55 + 555 + \dots$

$$\begin{aligned}
\text{Let } S_n &= 5 + 55 + 555 + \dots \text{ to } n \text{ terms} \\
&= \frac{5}{9} [9 + 99 + 999 + \dots \text{ to } n \text{ terms}] \\
&= \frac{5}{9} [(10 - 1) + (10^2 - 1) + (10^3 - 1) + \dots \text{ to } n \text{ terms}] \\
&= \frac{5}{9} [(10 + 10^2 + 10^3 + \dots n \text{ terms}) - (1 + 1 + \dots n \text{ terms})] \\
&= \frac{5}{9} \left[\frac{10(10^n - 1)}{10 - 1} - n \right] \\
&= \frac{5}{9} \left[\frac{10(10^n - 1)}{9} - n \right] \\
&= \frac{50}{81} (10^n - 1) - \frac{5n}{9}
\end{aligned}$$

(ii) .6 + .66 + .666 + ...

$$\begin{aligned}
\text{Let } S_n &= 0.6 + 0.66 + 0.666 + \dots \text{ to } n \text{ terms} \\
&= 6[0.1 + 0.11 + 0.111 + \dots \text{ to } n \text{ terms}] \\
&= \frac{6}{9} [0.9 + 0.99 + 0.999 + \dots \text{ to } n \text{ terms}] \\
&= \frac{6}{9} \left[\left(1 - \frac{1}{10}\right) + \left(1 - \frac{1}{10^2}\right) + \left(1 - \frac{1}{10^3}\right) + \dots \text{ to } n \text{ terms} \right] \\
&= \frac{2}{3} \left[(1 + 1 + \dots n \text{ terms}) - \frac{1}{10} \left(1 + \frac{1}{10} + \frac{1}{10^2} + \dots \text{ terms} \right) \right] \\
&= \frac{2}{3} \left[n - \frac{1}{10} \left(\frac{1 - \left(\frac{1}{10}\right)^n}{1 - \frac{1}{10}} \right) \right] \\
&= \frac{2}{3} n - \frac{2}{30} \times \frac{10}{9} (1 - 10^{-n}) \\
&= \frac{2}{3} n - \frac{2}{27} (1 - 10^{-n})
\end{aligned}$$

Page : 200 , Block Name : Miscellaneous Exercise

Q22 Find the 20th term of the series $2 \times 4 + 4 \times 6 + 6 \times 8 + \dots + n \text{ terms}$.

Answer.

The given series is $2 \times 4 + 4 \times 6 + 6 \times 8 + \dots$ n terms

$$\therefore n^{\text{th}} \text{ term} = a_n = 2n \times (2n + 2) = 4n^2 + 4n$$

$$a_{20} = 4(20)^2 + 4(20) = 4(400) + 80 = 1600 + 80 = 1680$$

Thus, the 20^{th} term of the series is 1680.

Page : 200 , Block Name : Miscellaneous Exercise

Q23 Find the sum of the first n terms of the series: $3 + 7 + 13 + 21 + 31 + \dots$

Answer.

The given series is $3 + 7 + 13 + 21 + 31 + \dots$

$$S = 3 + 7 + 13 + 21 + 31 + \dots + a_{n-1} + a_n$$

$$S = 3 + 7 + 13 + 21 + \dots + a_{n-2} + a_{n-1} + a_n$$

On subtracting both the equations, we obtain

$$S - S = [3 + (7 + 13 + 21 + 31 + \dots + a_{n-1} + a_n)] - [(3 + 7 + 13 + 21 + \dots + a_{n-2} + a_{n-1} + a_n)]$$

$$S - S = 3 + [(7 - 3) + (13 - 7) + (21 - 13) + \dots + (a_n - a_{n-1})]$$

$$0 = 3 + [4 + 6 + 8 + \dots (n - 1) \text{ terms}] - a_n$$

$$a_n = 3 + [4 + 6 + 8 + \dots (n - 1) \text{ terms}]$$

$$\begin{aligned} \Rightarrow a_n &= 3 + \left(\frac{n-1}{2} \right) [2 \times 4 + (n-1-1)2] \\ &= 3 + \left(\frac{n-1}{2} \right) [8 + (n-2)2] \end{aligned}$$

$$\begin{aligned}
&= 3 + \frac{(n-1)}{2}(2n+4) \\
&= 3 + (n-1)(n+2) \\
&= 3 + (n^2 + n - 2) \\
&= n^2 + n + 1 \\
\therefore \sum_{k=1}^n a_k &= \sum_{k=1}^n k^2 + \sum_{k=1}^n k + \sum_{k=1}^n 1 \\
&= \frac{n(n+1)(2n+1)}{6} + \frac{n(n+1)}{2} + n \\
&= n \left[\frac{(n+1)(2n+1) + 3(n+1) + 6}{6} \right] \\
&= n \left[\frac{2n^2 + 3n + 1 + 3n + 3 + 6}{6} \right] \\
&= n \left[\frac{2n^2 + 3n + 10}{6} \right] \\
&= \frac{n}{3} (n^2 + 3n + 5)
\end{aligned}$$

Page : 200 , Block Name : Miscellaneous Exercise

Q24

If S_1, S_2, S_3 are the sum of first n natural numbers, their squares cubes, respectively, show that $9S_2^2 = S_3 (1 + 8S_1)$

Answer. From the given information,

$$S_1 = \frac{n(n+1)}{2}$$

$$S_3 = \frac{n^2(n+1)^2}{4}$$

Here, $S_3 (1 + 8S_1) = \frac{n^2(n+1)^2}{4} \left[1 + \frac{8n(n+1)}{2} \right]$

$$= \frac{n^2(n+1)^2}{4} [1 + 4n^2 + 4n]$$

$$= \frac{n^2(n+1)^2}{4} (2n+1)^2$$

$$= \frac{[n(n+1)(2n+1)]^2}{4} \dots (1)$$

Also, $9S_2^2 = 9 \frac{[n(n+1)(2n+1)]^2}{4}$

$$= \frac{9}{36} [n(n+1)(2n+1)]^2$$

$$= \frac{[n(n+1)(2n+1)]^2}{4} \dots (2)$$

Thus, from (1) and (2), we obtain $9S_2^2 = S_3 (1 + 8S_1)$

Page : 200 , Block Name : Miscellaneous Exercise

Q25 Find the sum of the following series up to n terms:

$$\frac{1^3}{1} + \frac{1^3+2^3}{1+3} + \frac{1^3+2^3+3^3}{1+3+5} + \dots$$

Answer. The nth term of the given series is

$$\frac{1^3+2^3+3^3+\dots+n^3}{1+3+5+\dots+(2n-1)} = \frac{\left[\frac{n(n+1)}{2}\right]^2}{1+3+5+\dots+(2n-1)}$$

Here, 1, 3, 5, ..., (2n-1) is an A.P. with first term a, last term (2n-1) and number of terms as n

$$\therefore 1 + 3 + 5 + \dots + (2n-1) = \frac{n}{2} [2 \times 1 + (n-1)2] = n^2$$

$$\therefore a_n = \frac{n^2(n+1)^2}{4n^2} = \frac{(n+1)^2}{4} = \frac{1}{4}n^2 + \frac{1}{2}n + \frac{1}{4}$$

$$\therefore S_n = \sum_{k=1}^n a_k = \sum_{k=1}^n \left(\frac{1}{4}K^2 + \frac{1}{2}K + \frac{1}{4} \right)$$

$$= \frac{1}{4} \frac{n(n+1)(2n+1)}{6} + \frac{1}{2} \frac{n(n+1)}{2} + \frac{1}{4}n$$

$$= \frac{n[(n+1)(2n+1) + 6(n+1) + 6]}{24}$$

$$= \frac{n[2n^2 + 3n + 1 + 6n + 6 + 6]}{24}$$

$$= \frac{n(2n^2 + 9n + 13)}{24}$$

Page : 200 , Block Name : Miscellaneous Exercise

Q26 Show that $\frac{1 \times 2^2 + 2 \times 3^2 + \dots + n \times (n+1)^2}{1^2 \times 2 + 2^2 \times 3 + \dots + n^2 \times (n+1)} = \frac{3n+5}{3n+1}$.

Answer.

$$n^{\text{th}} \text{ term of the numerator} = n(n+1)^2 = n^3 + 2n^2 + n$$

$$n^{\text{th}} \text{ term of the denominator} = n^2(n+1) = n^3 + n^2$$

$$\frac{1 \times 2^2 + 2 \times 3^2 + \dots + n \times (n+1)^2}{1^2 \times 2 + 2^2 \times 3 + \dots + n^2 \times (n+1)} = \frac{\sum_{k=1}^n a_k}{\sum_{k=1}^n a_k} = \frac{\sum_{k=1}^n (K^3 + 2K^2 + K)}{\sum_{k=1}^n (K^3 + K^2)} \dots (1)$$

$$\begin{aligned} \text{Here, } \sum_{K=1}^n (K^3 + 2K^2 + K) &= \frac{n^2(n+1)^2}{4} + \frac{2n(n+1)(2n+1)}{6} + \frac{n(n+1)}{2} \\ &= \frac{n(n+1)[n(n+1)]}{2} + \frac{2}{3}(2n+1) + 1 \\ &= \frac{n(n+1)}{2} \left[\frac{3n^2 + 3n + 8n + 4 + 6}{6} \right] \\ &= \frac{n(n+1)}{12} [3n^2 + 11n + 10] \\ &= \frac{n(n+1)}{12} [3n^2 + 6n + 5n + 10] \\ &= \frac{n(n+1)}{12} [3n(n+2) + 5(n+2)] \\ &= \frac{n(n+1)(n+2)(3n+5)}{12} \dots (2) \end{aligned}$$

$$\begin{aligned} \text{Also, } \sum_{k=1}^n (K^3 + K^2) &= \frac{n^2(n+1)^2}{4} + \frac{n(n+1)(2n+1)}{6} \\ &= \frac{n(n+1)}{2} \left[\frac{n(n+1)}{2} + \frac{2n+1}{3} \right] \\ &= \frac{n(n+1)}{2} \left[\frac{3n^2 + 3n + 4n + 2}{6} \right] \\ &= \frac{n(n+1)}{12} [3n^2 + 7n + 2] \\ &= \frac{n(n+1)}{12} [3n^2 + 6n + n + 2] \\ &= \frac{n(n+1)}{12} [3n(n+2) + 1(n+2)] \\ &= \frac{n(n+1)(n+2)(3n+1)}{12} \dots (3) \end{aligned}$$

From (1), (2), and (3), we obtain

$$\begin{aligned} \frac{1 \times 2^2 + 2 \times 3^2 + \dots + n \times (n+1)^2}{1^2 \times 2 + 2^2 \times 3 + \dots + n^2 \times (n+1)} &= \frac{\frac{n(n+1)(n+2)(3n+5)}{12}}{\frac{n(n+1)(n+2)(3n+1)}{12}} \\ &= \frac{n(n+1)(n+2)(3n+5)}{n(n+1)(n+2)(3n+1)} = \frac{3n+5}{3n+1} \end{aligned}$$

Thus, the given result is proved.

Page : 200 , Block Name : Miscellaneous Exercise

Q27 A farmer buys a used tractor for Rs 12000.

He pays Rs 6000 cash and agrees to pay the balance in annual instalments of Rs 500 plus 12% interest on the unpaid amount.

How much will the tractor cost him?

Answer. It is given that the farmer pays Rs 6000 in cash.

Therefore, unpaid amount = Rs 12000 – Rs 6000 = Rs 6000

According to the given condition, the interest paid annually is

12% of 6000, 12% of 5500, 12% of 5000, ..., 12% of 500

Thus, total interest to be paid = 12% of 6000 + 12% of 5500 + 12% of 5000 + ... + 12% of 500

= 12% of (6000 + 5500 + 5000 + ... + 500)

= 12% of (500 + 1000 + 1500 + ... + 6000)

Now, the series 500, 1000, 1500 ... 6000 is an A.P. with both the first term and common difference equal to 500.

Let the number of terms of the A.P. be n.

$$\therefore 6000 = 500 + (n - 1) 500$$

$$\Rightarrow 1 + (n - 1) = 12$$

$$\Rightarrow n = 12$$

\therefore Sum of the A.P

$$= \frac{12}{2} [2(500) + (12 - 1)(500)] = 6[1000 + 5500] = 6(6500) = 39000$$

Thus, total interest to be paid = 12% of (500 + 1000 + 1500 + ... + 6000)

= 12% of 39000 = Rs 4680

Thus, cost of tractor = (Rs 12000 + Rs 4680) = Rs 16680

Page : 200 , Block Name : Miscellaneous Exercise

Q28 Shamshad Ali buys a scooter for Rs 22000.

He pays Rs 4000 cash and agrees to pay the balance in annual instalment of Rs 1000 plus 10% interest on the unpaid amount.

How much will the scooter cost him?

Answer. It is given that Shamshad Ali buys a scooter for Rs 22000 and pays Rs 4000 in cash.

\therefore Unpaid amount = Rs 22000 – Rs 4000 = Rs 18000

According to the given condition, the interest paid annually is

10% of 18000, 10% of 17000, 10% of 16000 ... 10% of 1000

Thus, total interest to be paid = 10% of 18000 + 10% of 17000 + 10% of 16000 + ... + 10% of 1000

= 10% of (18000 + 17000 + 16000 + ... + 1000)

= 10% of (1000 + 2000 + 3000 + ... + 18000)

Here, 1000, 2000, 3000 ... 18000 forms an A.P. with first term and common difference both equal to 1000.

Let the number of terms be n.

$$\therefore 18000 = 1000 + (n - 1) (1000)$$

$$\Rightarrow n = 18$$

$$\begin{aligned} \therefore 1000 + 2000 + \dots + 18000 &= \frac{18}{2} [2(1000) + (18 - 1)(1000)] \\ &= 9[2000 + 17000] \\ &= 171000 \end{aligned}$$

\therefore Total interest paid = 10% of (18000 + 17000 + 16000 + ... + 1000)

= 10% of Rs 171000 = Rs 17100

\therefore Cost of scooter = Rs 22000 + Rs 17100 = Rs 39100

Page : 200 , Block Name : Miscellaneous Exercise

Q29 A person writes a letter to four of his friends.

He asks each one of them to copy the letter and mail to four different persons with instruction that they move the chain similarly.

Assuming that the chain is not broken and that it costs 50 paise to mail one letter.

Find the amount spent on the postage when 8th set of letter is Mailed.

Answer. The numbers of letters mailed forms a G.P.: $4, 4^2, \dots, 4^8$

First term = 4

Common ratio = 4

Number of terms = 8

It is known that the sum of n terms of a G.P. is given by

$$S_n = \frac{a(r^n - 1)}{r - 1}$$

$$\therefore S_8 = \frac{4(4^8 - 1)}{4 - 1} = \frac{4(65536 - 1)}{3} = \frac{4(65535)}{3} = 4(21845) = 87380$$

It is given that the cost to mail one letter is 50 paise.

$$\therefore \text{Cost of mailing 87380 letters} = \text{Rs } 87380 \times \frac{50}{100} = \text{Rs } 43690$$

Thus, the amount spent when 8th set of letter is mailed is Rs 43690.

Page : 200 , Block Name : Miscellaneous Exercise

Q30 A man deposited Rs 10000 in a bank at the rate of 5% simple interest annually.

Find the amount in 15th year since he deposited the amount and also calculate the total amount after 20 years.

Answer. It is given that the man deposited Rs 10000 in a bank at the rate of 5% simple interest annually.

$$\therefore \text{Interest in first year} = \frac{5}{100} \times \text{Rs } 10000 = \text{Rs } 500$$

$$\therefore \text{Amount in 15th year} = \text{Rs } 10000 + \underbrace{500 + 500 + \dots + 500}_{14 \text{ times}}$$

$$= \text{Rs } 10000 + 14 \times \text{Rs } 500$$

$$= \text{Rs } 10000 + \text{Rs } 7000$$

$$= \text{Rs } 17000$$

$$\text{Amount after 20 years} = \text{Rs } 10000 + \underbrace{500 + 500 + \dots + 500}_{20 \text{ times}}$$

$$= \text{Rs } 10000 + 20 \times \text{Rs } 500$$

$$= \text{Rs } 10000 + \text{Rs } 10000$$

Page : 200 , Block Name : Miscellaneous Exercise

Q31 A manufacturer reckons that the value of a machine, which costs him Rs. 15625, will depreciate each year by 20%.

Find the estimated value at the end of 5 years.

Answer. Cost of machine = Rs 15625

Machine depreciates by 20% every year.

Therefore, its value after every year is 80% of the original cost i.e., $\frac{4}{5}$ of the original cost.

$$\therefore \text{Value at the end of 5 years} = 15625 \times \underbrace{\frac{4}{5} \times \frac{4}{5} \times \dots \times \frac{4}{5}}_{5 \text{ times}} = 5 \times 1024 = 5120$$